

Study on the Applicability of Varshni Potential to Predict the Mass Spectra of the Quark-Antiquark Systems in a Non-relativistic Framework

Etido P. Inyang^{a,c}, Ephraim P. Inyang^a, Eddy S. William^a and Etebong E. Ibekwe^b

^a Theoretical Physics Group, Department of Physics, University of Calabar, PMB 1115, Calabar, Nigeria.

^b Department of Physics, Akwa Ibom State University, Ikot Akpaden, P.M.B 1167, Uyo, Nigeria.

^c Department of Physics, National Open University of Nigeria, Jabi, Abuja

Doi: <https://doi.org/10.47011/14.4.8>

Received on: 14/10/2020;

Accepted on: 29/11/2020

Abstract: In this work, we obtain the Schrödinger equation solutions for the Varshni potential using the Nikiforov-Uvarov method. The energy eigenvalues are obtained in non-relativistic regime. The corresponding eigenfunction is obtained in terms of Laguerre polynomials. We applied the present results to calculate heavy-meson masses of charmonium $c\bar{c}$ and bottomonium $b\bar{b}$. The mass spectra for charmonium and bottomonium multiplets have been predicted numerically. The results are in good agreement with experimental data and the works of other researchers.

Keywords: Schrödinger equation, Varshni potential, Nikiforov-Uvarov method, Heavy meson.

PACs: 14.20.Lq; 03.65.-w; 14.40.Pq; 11.80.Fv.

1. Introduction

It is a well-known fact by many researchers that quarkonium systems provide deep insight into the essential description of quantum chromodynamics (QCD) theory, particle physics and standard model [1-3]. Quarkonia with heavy quark and antiquark and their interaction are well described by the Schrödinger equation (SE). The solution of this SE with potential is one of the most fundamental problems in quarkonium systems. It is noted that the potentials considered should take into account two important features of the strong interaction; namely, asymptotic freedom and quark confinement [4-8]. The most fundamental potential used in studying quarkonium system is the Cornell potential, also known as Killingbeck potential. Most researchers have carried out works with Cornell potential [9, 10]. For instance, Vega and Flores

[11] solved the Schrödinger equation with the Cornell potential using the variational method and supersymmetric quantum mechanics (SUSYQM). Ciftci and Kisoglu [12] addressed non-relativistic arbitrary l -states of quark-antiquark through the Asymptotic Iteration Method (AIM). The energy eigenvalues with any $l \neq 0$ states and mass of the massive quark-antiquark system (quarkonium) were gotten. An analytic solution of the N-dimensional radial Schrödinger equation with the mixture of vector and scalar potentials *via* the Laplace transformation method (LTM) was studied by [13]. Their results were employed to analyze the different properties of the heavy-light mesons. Al-Jamel and Widyan [14] studied heavy quarkonium mass spectra in a Coulomb field plus quadratic potential using the Nikiforov-

Uvarov method. In their work, the spin-averaged mass spectra of heavy quarkonia in a Coulomb plus quadratic potential is analyzed within the non-relativistic Schrödinger equation. Al-Oun *et al.* [15] examined heavy quarkonia characteristics in the general framework of a non-relativistic potential model consisting of a Coulomb plus quadratic potential. Kumar and Chand [16] carried out an asymptotic study to the N-dimensional radial Schrödinger equation for the quark-antiquark interaction potential employing asymptotic iteration method (AIM). Ibekwe *et al.* [17] solved the radial SE with an exponential, generalized, anharmonic Cornell potential using the series expansion method. They applied the bound state eigenvalues to study the energy spectra for CO, NO, CH and N₂ diatomic molecules and the mass spectra of heavy quarkonium systems. Furthermore, Omugbe *et al.* [18] solved the SE with Killingbeck potential plus an inversely quadratic potential model. They obtained the energy eigenvalues and the mass spectra of the heavy and heavy-light meson systems. In addition, Ali *et al.* [19] studied the energy spectra of mesons and hadronic interactions using Numerov's method. Their solutions were used to describe the phenomenological interactions between the charm-anticharm quarks *via* the model. The model accurately predicts the mass spectra of charmed quarkonium as an example of mesonic

systems. Inyang *et al.* [20] obtained the Klein-Gordon equation solutions for the Yukawa potential using the Nikiforov-Uvarov method. The energy eigenvalues were obtained both in relativistic and non-relativistic regime. They applied the results to calculate heavy-meson masses of charmonium and bottomonium.

The Varshni potential is greatly important with applications, cutting across nuclear physics, particle physics and molecular physics [21]. The Varshni potential takes the form:

$$V(r) = a - \frac{abe^{-\alpha r}}{r} \quad (1)$$

where a and b are potential strengths for Varshni potential, α is the screening parameter which controls the shape of the potential energy curve as shown in Fig. 1 and r is the inter-nuclear separation. The Varshni potential is a short-range repulsive potential energy that plays an important role in chemical, particle and molecular physics [22]. This potential is used generally to describe bound states of the interaction of systems and has been applied in both classical and molecular physics. The Varshni potential was studied by Lim using the 2-body Kaxiras-Pandey parameters. He observed that Kaxiras and Pandey used this potential to describe the 2-body energy portion of multi-body condensed matter [23].




FIG.1. Plots of Varshni potential with r in (fm^{-1}).

Many researchers have studied heavy mesons with non-exponential type potential like the Cornell without considering the exponential type [11-16]. Therefore, we intend to investigate the SE with the exponential type potential (Varshni potential) in the framework of NU method to obtain the mass spectra of quark- antiquark systems. To the best of our knowledge, this is the first time Varshni potential is being studied with the aim of determining the mass spectra of heavy quarkonia systems using the NU method.

The paper is organized as follows: In Section 2, the Nikiforov-Uvarov (NU) method is reviewed. In Section 3, the bound state energy eigenvalues and the corresponding eigenfunction are calculated. In Section 4, the results are discussed. In Section 5, the conclusion is presented.

2. A Brief Review of Nikiforov-Uvarov (NU) Method

The NU method is used to transform Schrödinger -like equations into a second-order differential equation through a coordinate transformation $y = y(r)$, of the form [24]:

$$\psi''(y) + \frac{\tilde{\tau}(y)}{\sigma(y)}\psi'(y) + \frac{\tilde{\sigma}(y)}{\sigma^2(y)}\psi(y) = 0 \quad (2)$$

where $\tilde{\sigma}(y)$ and $\sigma(y)$ are polynomials, at most second-degree and $\tilde{\tau}(y)$ is a first-degree polynomial. From Eq. (2), we obtain exact solution by using the transformation:

$$\psi(y) = \phi(y)\chi(y). \quad (3)$$

This transformation reduces Eq. (2) into a hypergeometric-type equation of the form:

$$\sigma(y)\chi''(y) + \tau(y)\chi'(y) + \lambda\chi(y) = 0. \quad (4)$$

The function $\phi(y)$ can be defined as the logarithm derivative:

$$\frac{\phi'(y)}{\phi(y)} = \frac{\pi(y)}{\sigma(y)} \quad (5)$$

where $\pi(y)$ is at most a first-degree polynomial. The second part of the wave functions in Eq. (4) is a hypergeometric-type function obtained by Rodriguez relation:

$$\chi_n(y) = \frac{B_n}{\rho(y)} \frac{d^n}{dy^n} [\sigma^n(y)\rho(y)] \quad (6)$$

where B_n is the normalization constant and $\rho(y)$ the weight function which satisfies the condition below:

$$\frac{d}{dy}(\sigma(y)\rho(y)) = \tau(y)\rho(y) \quad (7)$$

where also:

$$\tau(y) = \tilde{\tau}(y) + 2\pi(y). \quad (8)$$

For bound solutions, it is required that:

$$\frac{d\tau(y)}{dy} < 0. \quad (9)$$

With $\pi(y)$ and parameter λ , the eigenfunctions and eigenvalues can be obtained using the definition of the following function:

$$\pi(y) = \frac{\sigma'(y) - \tilde{\tau}(y)}{2} \pm \sqrt{\left(\frac{\sigma'(y) - \tilde{\tau}(y)}{2}\right)^2 - \tilde{\sigma}(y) + k\sigma(y)} \quad (10)$$

and

$$\lambda = k + \pi'(y). \quad (11)$$

The value of k can be obtained by setting the discriminant in the square root in Eq. (10) equal to zero. As such, the new eigenvalues equation can be given as:

$$\lambda + n\tau'(y) + \frac{n(n-1)}{2}\sigma''(y) = 0, (n = 0, 1, 2, \dots). \quad (12)$$

3. Approximate Solutions of the Schrödinger Equation with Varshni Potential

The Schrödinger equation (SE) for two particles interacting *via* potential $V(r)$ in three-dimensional space is given by [25]:

$$\frac{d^2 R(r)}{dr^2} + \left[\frac{2\mu}{\hbar^2} (E_n - V(r)) - \frac{l(l+1)}{r^2} \right] R(r) = 0 \quad (13)$$

where l, μ, r and \hbar are the angular momentum quantum number, the reduced mass for the quarkonium particle, inter-particle distance and reduced plank constant, respectively.

We carry out Taylor series expansion of the exponential term in Eq. (1) up to order three, in order to make the potential interact in the quark-antiquark system and this yields:

$$\frac{e^{-\alpha r}}{r} = \frac{1}{r} - \alpha + \frac{\alpha^2 r}{2} - \frac{\alpha^3 r^2}{6} + \dots \quad (14)$$

We substitute Eq. (14) into Eq. (1) and obtain:

$$V(r) = -\frac{B}{r} - Cr + Dr^2 + A \quad (15)$$

where

$$\left. \begin{aligned} A &= a + ab\alpha, B = ab \\ C &= \frac{ab\alpha^2}{2}, D = \frac{ab\alpha^3}{6} \end{aligned} \right\}. \quad (16)$$

Upon substituting Eq. (15) into Eq. (13), we obtain:

$$\frac{d^2 R(r)}{dr^2} + \left[\frac{2\mu E_{nl}}{\hbar^2} + \frac{2\mu B}{\hbar^2 r} + \frac{2\mu Cr}{\hbar^2} - \frac{2\mu Dr^2}{\hbar^2} - \frac{2\mu A}{\hbar^2} - \frac{l(l+1)}{r^2} \right] R(r) = 0. \quad (17)$$

In order to transform the coordinate from r to y in Eq. (17), we set:

$$y = \frac{1}{r}. \quad (18)$$

This implies that the 2nd derivative in Eq. (18) becomes:

$$\frac{d^2 R(r)}{dr^2} = 2y^3 \frac{dR(y)}{dy} + y^4 \frac{d^2 R(y)}{dy^2}. \quad (19)$$

Substituting Eqs. (18) and (19) into Eq. (17), we obtain:

$$\frac{d^2 R(y)}{dy^2} + \frac{2y}{y^2} \frac{dR(y)}{dy} + \frac{1}{y^4} \left[\frac{2\mu E_{nl}}{\hbar^2} + \frac{2\mu By}{\hbar^2} + \frac{2\mu C}{\hbar^2 y} - \frac{2\mu D}{\hbar^2 y^2} - \frac{2\mu A}{\hbar^2} - l(l+1)y^2 \right] R(y) = 0. \quad (20)$$

Next, we propose the following approximation scheme on the term $\frac{C}{y}$ and $\frac{D}{y^2}$.

Let us assume that there is a characteristic radius r_0 of the meson. Then, the scheme is based on the expansion of $\frac{C}{y}$ and $\frac{D}{y^2}$ in a power series around r_0 ; i.e., around $\delta \equiv \frac{1}{r_0}$, up to the second order. This is similar to Pekeris approximation, which helps deform the centrifugal term such that the modified potential can be solved by the NU method [26].

Setting $x = y - \delta$ and around $x = 0$, it can be expanded into a series of powers as:

$$\frac{C}{y} = \frac{C}{x + \delta} = \frac{C}{\delta \left(1 + \frac{x}{\delta} \right)} = \frac{C}{\delta} \left(1 + \frac{x}{\delta} \right)^{-1} \quad (21)$$

which yields:

$$\frac{C}{y} = C \left(\frac{3}{\delta} - \frac{3y}{\delta^2} + \frac{y^2}{\delta^3} \right). \quad (22)$$

Similarly,

$$\frac{D}{y^2} = D \left(\frac{6}{\delta^2} - \frac{8y}{\delta^3} + \frac{3y^2}{\delta^4} \right). \quad (23)$$

Substituting Eqs. (22) and (23) into Eq.(20) yields:

$$\frac{d^2 R(y)}{dy^2} + \frac{2y}{y^2} \frac{dR(y)}{dy} + \frac{1}{y^4} [-\varepsilon + \alpha y - \beta y^2] R(y) = 0 \quad (24)$$

where

$$\left. \begin{aligned} -\varepsilon &= \left(\frac{2\mu E_{nl}}{\hbar^2} - \frac{2\mu A}{\hbar^2} + \frac{6\mu C}{\hbar^2 \delta} - \frac{12\mu D}{\hbar^2 \delta^2} \right), \\ \alpha &= \left(\frac{2\mu B}{\hbar^2} - \frac{2\mu C}{\hbar^2 \delta^2} + \frac{16\mu D}{\hbar^2 \delta^3} \right) \\ \beta &= \left(\gamma - \frac{2\mu C}{\hbar^2 \delta^3} + \frac{6\mu D}{\hbar^2 \delta^4} \right), \gamma = l(l+1) \end{aligned} \right\}. \quad (25)$$

Comparing Eq. (24) and Eq.(2), we obtain:

$$\left. \begin{aligned} \tilde{\tau}(y) &= 2y, \sigma(y) = y^2, \\ \tilde{\sigma}(y) &= -\varepsilon + \alpha y - \beta y^2 \\ \sigma'(y) &= 2y, \sigma''(y) = 2 \end{aligned} \right\}. \quad (26)$$

Substituting Eq. (26) into Eq. (10) yields:

$$\pi(y) = \pm \sqrt{\varepsilon - \alpha y + (\beta + k)y^2}. \quad (27)$$

To determine k , we take the discriminant of the function under the square root, which yields:

$$k = \frac{\alpha^2 - 4\beta\varepsilon}{4\varepsilon}. \quad (28)$$

We substitute Eq. (28) into Eq. (27) and have:

$$\pi(y) = \pm \left(\frac{\alpha y}{2\sqrt{\varepsilon}} - \frac{\varepsilon}{\sqrt{\varepsilon}} \right). \quad (29)$$

We take the negative part of Eq. (29) and differentiate, which yields:

$$\pi'_-(y) = -\frac{\alpha}{2\sqrt{\varepsilon}}. \quad (30)$$

By substituting Eqs. (26) and (29) into Eq.(8), we have:

$$\tau(y) = 2y - \frac{\alpha y}{\sqrt{\varepsilon}} + \frac{2\varepsilon}{\sqrt{\varepsilon}}. \quad (31)$$

Differentiating Eq. (31), we have:

$$\tau'(y) = 2 - \frac{\alpha}{\sqrt{\varepsilon}}. \quad (32)$$

By using Eq. (11), we obtain:

$$\lambda = \frac{\alpha^2 - 4\beta\epsilon}{4\epsilon} - \frac{\alpha}{2\sqrt{\epsilon}}. \quad (33)$$

And using Eq. (12), we obtain:

$$\lambda_n = \frac{n\alpha}{\sqrt{\epsilon}} - n^2 - n. \quad (34)$$

Equating Eqs. (33) and (34), the energy eigenvalue of Eq. (17) is given:

$$E_{nl} = a(1 + b\alpha) - \frac{3ab\alpha^3}{2\delta} + \frac{ab\alpha^3}{\delta^3} - \frac{\hbar^2}{8\mu} \left[\frac{\frac{2\mu ab}{\hbar^2} - \frac{3\mu ab\alpha^2}{\hbar^2\delta^2} + \frac{16\mu ab\alpha^3}{6\hbar^2\delta^2}}{n + \frac{1}{2} + \sqrt{\left(l + \frac{1}{2}\right)^2 - \frac{ab\mu\alpha^3}{\delta^3\hbar^2} + \frac{16\mu ab\alpha^3}{6\hbar^2\delta^4}}} \right]^2. \quad (35)$$

To determine the wavefunction, we substitute Eqs. (26) and (29) into Eq.(5) and obtain:

$$\frac{d\phi}{\phi} = \left(\frac{\epsilon}{y^2\sqrt{\epsilon}} - \frac{\alpha}{2y\sqrt{\epsilon}} \right) dy. \quad (36)$$

Integrating Eq. (36) gives:

$$\phi(y) = y^{-\frac{\alpha}{2\sqrt{\epsilon}}} e^{-\frac{\epsilon}{y\sqrt{\epsilon}}}. \quad (37)$$

By substituting Eqs. (26) and (29) into Eq. (7), integrating and thereafter simplifying, we obtain:

$$\rho(y) = y^{-\frac{\alpha}{\sqrt{\epsilon}}} e^{-\frac{2\epsilon}{y\sqrt{\epsilon}}}. \quad (38)$$

Substituting Eqs. (26) and (38) into Eq.(6), we have:

$$\chi_n(y) = B_n e^{\frac{2\epsilon}{y\sqrt{\epsilon}}} y^{\frac{\alpha}{\sqrt{\epsilon}}} \frac{d^n}{dy^n} \left[e^{-\frac{2\epsilon}{y\sqrt{\epsilon}}} y^{2n - \frac{\alpha}{\sqrt{\epsilon}}} \right]. \quad (39)$$

The Rodrigues' formula of the associated Laguerre polynomials is:

$$L_n^{\frac{\alpha}{\sqrt{\epsilon}}} \left(\frac{2\epsilon}{y\sqrt{\epsilon}} \right) = \frac{1}{n!} e^{\frac{2\epsilon}{y\sqrt{\epsilon}}} y^{\frac{\alpha}{\sqrt{\epsilon}}} \frac{d^n}{dy^n} \left(e^{-\frac{2\epsilon}{y\sqrt{\epsilon}}} y^{2n - \frac{\alpha}{\sqrt{\epsilon}}} \right) \quad (40)$$

where

$$\frac{1}{n!} = B_n. \quad (41)$$

Hence,

$$\chi_n(y) \equiv L_n^{\frac{\alpha}{\sqrt{\epsilon}}} \left(\frac{2\epsilon}{y\sqrt{\epsilon}} \right). \quad (42)$$

Substituting Eqs. (37) and (42) into Eq. (3), we obtain the wavefunction of Eq.(17) in terms of Laguerre polynomials as:

$$\psi(y) = N_{nl} y^{-\frac{\alpha}{2\sqrt{\epsilon}}} e^{-\frac{\epsilon}{y\sqrt{\epsilon}}} L_n^{\frac{\alpha}{\sqrt{\epsilon}}} \left(\frac{2\epsilon}{y\sqrt{\epsilon}} \right) \quad (43)$$

where N_{nl} is normalization constant, which can be obtained from:

$$\int_0^\infty |N_{nl}(r)|^2 dr = 1. \quad (44)$$

4. Results and Discussion

4.1 Results

The mass spectra of the heavy quarkonium system such as charmonium and bottomonium that have the quark and antiquark flavor are calculated and we apply the following relation: [27, 28]

$$M = 2m + E_{nl}, \quad (45)$$

where m is quarkonium bare mass and E_{nl} is energy eigenvalue. By substituting Eq. (35) into Eq. (45), we obtain the mass spectra for Varshni potential as:

$$M = 2m + a(1 + b\alpha) - \frac{3ab\alpha^3}{2\delta} + \frac{ab\alpha^3}{\delta^3} - \frac{\hbar^2}{8\mu} \left[\frac{\frac{2\mu ab}{\hbar^2} - \frac{3\mu ab\alpha^2}{\hbar^2\delta^2} + \frac{16\mu ab\alpha^3}{6\hbar^2\delta^2}}{n + \frac{1}{2} + \sqrt{\left(l + \frac{1}{2}\right)^2 - \frac{ab\mu\alpha^3}{\delta^3\hbar^2} + \frac{16\mu ab\alpha^3}{6\hbar^2\delta^4}}} \right]^2. \quad (46)$$

In order to test for the accuracy of the predicted results determined numerically, we used a Chi square function defined by [29]:

$$\chi^2 = \frac{1}{n} \sum_{i=1}^n \frac{(M_i^{\text{Exp.}} - M_i^{\text{Theo.}})^2}{\Delta_i} \quad (47)$$

where n runs over selected samples of heavy mesons, $M_i^{\text{Exp.}}$ is the experimental mass of heavy-mesons, while M_i^{Th} is the corresponding theoretical prediction. The Δ_i quantity is experimental uncertainty of the masses. Intuitively, Δ_i should be one. The tendency of overestimating Chi square value is that it reflects some mean error per heavy meson state.

4.2 Discussion of Results

We calculate mass spectra of charmonium and bottomonium for states from 1S to 1F by using Eq. (46). We adopt the numerical values of bottomonium ($b\bar{b}$) and charmonium ($c\bar{c}$) masses as 4.68 GeV and 1.488 GeV , respectively, Ref. [30]. Then, the corresponding reduced masses are $\mu_b = 2.340 \text{ GeV}$ and $\mu_c = 0.744 \text{ GeV}$. The free parameters of Eq.(46) were then obtained by solving two algebraic equations by inserting experimental data of mass spectra for $2S, 2P$ in the case of charmonium. In the case of bottomonium, the values of the free parameters in Eq. (46) are calculated by solving two algebraic equations, which were obtained by inserting experimental data of mass spectra for $1S, 2S$. Experimental data is taken from Ref. [31].

We note that calculations of mass spectra of charmonium and bottomonium are in good agreement with the experimental data, given that the maximum error in comparison with the

experimental data is 0.0055 GeV . The values obtained are in good agreement with the works of other researchers, like Ref.[24] and Ref.[18], as shown in Tables 1 and 2. In Ref. [24], the authors investigated the N-radial SE analytically by employing Cornell potential, which was extended to finite temperature. In Ref.[18], the Klein-Gordon equation is solved for the Yukawa potential using the Nikiforov-Uvarov method. The energy eigenvalues were obtained both in relativistic and non-relativistic regime. The results were used to calculate heavy-meson masses of charmonium $c\bar{c}$ and bottomonium $b\bar{b}$. We also plotted mass spectra energy against potential strength (a), reduced mass (μ) and screening parameter (α), respectively. In Fig. 2, the mass spectra energy converge at the beginning but spread out and there is a monotonic increase in potential strength (a). Figs. 3 and 4 show the convergence of the mass spectra energy as the screening parameter (α) and reduced mass (μ) increase for various angular momentum quantum numbers. This indicates the energy peak as observable to determine the top quark mass.

TABLE 1. Mass spectra of charmonium in (GeV) ($m_c = 1.488 \text{ GeV}$, $\mu = 0.744 \text{ GeV}$, $\alpha = -0.976$, $\delta = 1.700 \text{ GeV}$, $\hbar = 1$, $a = -48.049 \text{ GeV}$ and $b = 3.020 \text{ GeV}$).

State	Present work	[26]	[20]	Experiment[31]
1S	3.096	3.096	3.096	3.096
2S	3.686	3.686	3.672	3.686
1P	3.295	3.255	3.521	3.525
2P	3.802	3.779	3.951	3.773
3S	4.040	4.040	4.085	4.040
4S	4.269	4.269	4.433	4.263
1D	3.583	3.504	3.800	3.770
2D	3.976	-	-	4.159
1F	3.862	-	-	-

TABLE 2. Mass spectra of bottomonium in (GeV) ($m_b = 4.680 \text{ GeV}$, $\mu = 2.340 \text{ GeV}$, $\alpha = -0.952$, $\delta = 1.70 \text{ GeV}$, $\hbar = 1$, $a = -14.352 \text{ GeV}$ and $b = 3.084 \text{ GeV}$).

State	Present work	[26]	[20]	Experiment[31]
1S	9.460	9.460	9.462	9.460
2S	10.569	10.023	10.027	10.023
1P	9.661	9.619	9.963	9.899
2P	10.138	10.114	10.299	10.260
3S	10.355	10.355	10.361	10.355
4S	10.567	10.567	10.624	10.580
1D	9.943	9.864	10.209	10.164
2D	10.306	-	-	-
1F	10.209	-	-	-




FIG. 2. Variation of mass spectra with potential strength (a) for different quantum numbers.




FIG. 3. Variation of mass spectra with reduced mass μ for different quantum numbers.




FIG. 4. Variation of mass spectra with screening parameter (α) for different quantum numbers.

5. Conclusion

In this work, the bound-state solutions of the Schrödinger equation for the Varshni potential using the Nikiforov-Uvarov method were obtained. The corresponding eigenfunction was achieved in terms of Laguerre polynomials. We applied the present results to calculate heavy-meson masses such as charmonium and bottomonium. The energy eigenvalues of charmonium ($c\bar{c}$) and bottomonium ($b\bar{b}$) for

states 1S to 1F were obtained and compared with experimental data and other theoretical works, which are in good agreement with the maximum error of 0.0055 GeV . The exponential type potential has been successfully applied in predicting the mass spectra of heavy mesons. The analytical solutions can also be used to describe other characteristics of the quarkonium systems like thermodynamic properties.

References

- [1] Abu-Shady, M., Abdel-Karim, T.A. and Khokha E.M., Adv. High-ener. Phys., 43 (2018) 843.
- [2] Bhanot, G. and Rudaz, S., Phys. Lett. B, 78 (1978) 119.
- [3] Ntibi, J.E., Inyang, E.P., Inyang, E.P. and William, E.S., Int. J. Innova. Sci. Eng. Tech., 11 (2020) 2348.
- [4] Martin, B.R. and Shaw, G.G., "Particle Physics", 2nd edition, (The Manchester Physics Series, 1997).
- [5] Perkins, D.H., "Introduction to High-energy Physics", (Cambridge University Press, Cambridge, UK, 20, 2000).
- [6] Griffiths, D., "Introduction to Elementary Particles", (John Wiley & Sons, New York, USA, 1987).
- [7] Bettini, A., "Introduction to Elementary Particle Physics", (Cambridge University Press, 2018).
- [8] Rathsman, J., "Quark and Lepton Interactions", (Acta Universitatis Upsalensis Uppsala, 1996).

-
- [9] Abu-Shady, M., *Int. J. Appl. Math. and Theor. Phys.*, 2 (2) (2016) 16.
- [10] Omugbe, E., *Cana. J. Phys.*, 98 (12) (2020) 1125.
- [11] Vega, A. and Flores, J., *Pramana J. Phys.*, 56 (2016) 467.
- [12] Ciftci, H. and Kisoglu, H.F., *Pramana J. Phys.*, 57 (2018) 467.
- [13] Abu-Shady, M., *Bos. J. Mod. Phys.*, 55 (2015) 799.
- [14] Al-Jamel, A. and Widyan, H., *Cana. Cent. Sci. Educ.*, 4 (2012) 29.
- [15] Al-Oun, A., Al-Jamel, A. and Widyan, H., *Jordan. J. Phys.*, 8 (4) (2015) 199.
- [16] Kumar, R. and Chand, F., *Comm. Theor. Phys.*, 59 (2013) 467.
- [17] Ibekwe, E.E., Alalibo, T.N., Uduakobong, S.O., Ikot, A.N. and Abdullah, N.Y., *Iran J. Sci. Technol. Trans. Sci.*, 20 (2020) 913.
- [18] Omuge, E., Osafire, O.E. and Onyeajh, M.C., *Adv. High-ener. Phys.*, 10 (2020) 1155.
- [19] Ali, M.S., Hassan, G.S., Abdelmonem, A.M., Elshamndy, S.K., Elmaety, F. and Yasser, A.M., *J. Rad. Appl. Sci.*, 13 (2020) 233.
- [20] Inyang, E.P., Inyang, E.P., Ntibi, J.E., Ibekwe, E.E. and William, E.S., *Indian J. of Phys.*, 1 (2021) 0974.
- [21] Varshni, Y.P., *Rev. Mod. Phys.*, 29 (1957) 682.
- [22] Oluwadare, O.J. and Oyewumi, K.J., *Eur. Phys.*, 53 (2017) 296.
- [23] Lim, T.C., *J. Serb. Chem. Soc.*, 74 (2009) 1423.
- [24] Nikiforov, A.F. and Uvarov, V.B., "Mat. Phys." (ed.) A. Jaffe (Germany: Birkhauser Verlag Basel, 317, 1988).
- [25] William, E.S., Inyang, E.P. and Thompson, E.A., *Rev. Mex. de Fis.*, 66 (6) (2020) 741.
- [26] Abu-Shady, M., *J. Egyp. Math. Soc.*, 23 (2016) 4.
- [27] Kuchin, S.M. and Maksimenko, N.V., *Univ. J. Phys. Appl.*, 7 (2013) 295.
- [28] Abu-Shady, M., Abdel-Karim, T.A. and Khokha, E.M., *J. Quan. Phys.*, 2 (2018) 17.
- [29] Ali, M.S., Yasser, A.M., Hassan, G.S. and Moustakidis, C.C., *Quan. Phys. Lett.*, 5 (2016) 14.
- [30] Barnett, R.M., Carone, C.D., Groom, D.E., Trippe, T.G. and Wohl, C.G., *Phys. Rev. D*, 92 (2012) 656.
- [31] Tanabashi, M., Carone, C.D., Trippe, T.G. and Wohl, C.G., *Phys. Rev. D*, 98 (2018) 548.