

# Jordan Journal of Physics

## ARTICLE

---

### About the Cosmological Constant, Acceleration Field, Pressure Field and Energy

**Sergey G. Fedosin**

*Sviazeva Str. 22-79, Perm, 614088, Russia.*

---

*Received on: 12/9/2015;*

*Accepted on: 26/11/2015*

---

**Abstract:** Based on the condition of relativistic energy uniqueness, the calibration of the cosmological constant was performed. This allowed to obtain the corresponding equation for the metric and to determine the generalized momentum, the relativistic energy, momentum and mass of the system, as well as the expressions for the kinetic and potential energies. The scalar curvature at an arbitrary point of the system equaled zero, if the matter is absent at this point; the presence of a gravitational or electromagnetic field is enough for the space-time curvature. Four-potentials of the acceleration field and pressure field, as well as tensor invariants determining the energy density of these fields, were introduced into the Lagrangian in order to describe the system's motion more precisely. The structure of the Lagrangian used is completely symmetrical in form with respect to the four-potentials of gravitational, electromagnetic, acceleration and pressure fields. The stress-energy tensors of the gravitational, acceleration and pressure fields are obtained in explicit form. Each of them can be expressed through the corresponding field vector and additional solenoidal vector. A description of the equations of acceleration and pressure fields is provided.

**Keywords:** Cosmological constant, Acceleration field, Pressure field, Covariant theory of gravitation.

## 1. Introduction

The most popular application of the cosmological constant  $\Lambda$  in the general theory of relativity (GTR) is that this quantity represents the manifestation of the vacuum energy [1-2]. There is another approach to the cosmological constant interpretation, according to which this quantity represents the energy possessed by any solitary particle in the absence of external fields. In this case, including  $\Lambda$  into the Lagrangian seems quite appropriate, since the Lagrangian contains such energy components, which should fully describe the properties of any system consisting of particles and fields.

Earlier in [3-4], we used such calibration of the cosmological constant, which allowed to maximally simplify the equation for the metric.

The disadvantage of this approach was that the relativistic energy of the system could not be determined uniquely, since the expression for the energy included the scalar curvature. In this paper, we use another universal calibration of the cosmological constant, which is suitable for any particle and system of particles and fields. As a result, the energy is independent of both the scalar curvature and the cosmological constant.

In GTR, the gravitational field as a separate object is not included in the Lagrangian and the role of a field is played by the metric itself. A known problem arising from such an approach is that in GTR there is no stress-energy tensor of the gravitational field.

In contrast, in the covariant theory of gravitation, the Lagrangian is used containing

the term with the energy of the particles in the gravitational field and the term with the energy of the gravitational field as such. Thus, the gravitational field is included in the Lagrangian in the same way as the electromagnetic field. In this case, the metric of the curved spacetime is used to specify the equations of motion as compared to the case of such a weak field, the limit of which is the special theory of relativity. In the weak field limit, a simplified metric is used, which almost does not depend on the coordinates and time. This is enough in many cases, for example, in case of describing the motion of planets. However, generally, in case of strong fields and for studying the subtle effects, the use of metric becomes necessary.

We will note that the term with the particle energy in the Lagrangian can be written in different ways. In [5-6], this term contains the invariant  $c\rho_0\sqrt{g_{\mu\nu}u^\mu u^\nu}$ , where  $\rho_0$  is the mass density in the co-moving reference frame and  $u^\mu$  is four-velocity. The corresponding quantity in [7] has the form  $\rho_0 c^2$ . In [3] and [8], instead of it the product  $c\sqrt{g_{\mu\nu}J^\mu J^\nu}$  is used, where

$$S = \int L dt = \int \left( k(R - 2\Lambda) - \frac{1}{c} D_\mu J^\mu + \frac{c}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{c} A_\mu j^\mu - \frac{c\varepsilon_0}{4} F_{\mu\nu} F^{\mu\nu} - \right. \\ \left. - \frac{1}{c} u_\mu J^\mu - \frac{c}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{1}{c} \pi_\mu J^\mu - \frac{c}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} d\Sigma, \quad (1)$$

where:

- $L$  – the Lagrange function or Lagrangian,
- $dt$  – the differential of the coordinate time of the used reference frame,
- $k$  – a coefficient to be determined,
- $R$  – the scalar curvature,
- $\Lambda$  – the cosmological constant,
- $J^\mu = \rho_0 u^\mu$  – the four-vector of gravitational (mass) current,
- $\rho_0$  – the mass density in the reference frame associated with the particle,
- $u^\mu = \frac{c dx^\mu}{ds}$  – the four-velocity of a point particle,  $dx^\mu$  – four-displacement,  $ds$  – interval,

$J^\mu$  is mass four-current. In this paper, we have chosen another form of the mentioned invariant – in the form  $u_\mu J^\mu$ . The reason for this choice is the fact that we consider the mass four-current  $J^\mu = \rho_0 u^\mu$  to be the fullest representative of the properties of matter particles containing both the mass density and the four-velocity. The mass four-current can be considered as the four-potential of the matter field. All the other four-vectors in the Lagrangian are four-potentials of the respective fields and are written with covariant indices. With the help of these four-potentials, tensor invariants are calculated which characterize the energy of the respective field in the Lagrangian.

## Action and its variations in the principle of least action

### The action function

We use the following expression as the action function for continuously distributed matter in the gravitational and electromagnetic fields in an arbitrary frame of reference:

$c$  – the speed of light as a measure of the propagation velocity of electromagnetic and gravitational interactions,

$D_\mu = \left( \frac{\psi}{c}, -\mathbf{D} \right)$  – the four-potential of the gravitational field, described by the scalar potential  $\psi$  and the vector potential  $\mathbf{D}$  of this field,

$G$  – the gravitational constant,

$\Phi_{\mu\nu} = \nabla_\mu D_\nu - \nabla_\nu D_\mu = \partial_\mu D_\nu - \partial_\nu D_\mu$  – the gravitational tensor (the tensor of gravitational field strengths),

$\Phi^{\alpha\beta} = g^{\alpha\mu} g^{\nu\beta} \Phi_{\mu\nu}$  – definition of the gravitational tensor with contravariant indices by means of the metric tensor  $g^{\alpha\mu}$ ,

$A_\mu = \left( \frac{\varphi}{c}, -\mathbf{A} \right)$  – the four-potential of the electromagnetic field, which is set by the scalar potential  $\varphi$  and the vector potential  $\mathbf{A}$  of this field,

$j^\mu = \rho_{0q} u^\mu$  – the four-vector of the electromagnetic (charge) current,

$\rho_{0q}$  – the charge density in the reference frame associated with the particle,

$\varepsilon_0$  – the vacuum permittivity,

$F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu = \partial_\mu A_\nu - \partial_\nu A_\mu$  – the electromagnetic tensor (the tensor of electromagnetic field strengths),

$u_\mu = g_{\mu\nu} u^\nu$  – the four-velocity with a covariant index, expressed through the metric tensor and the four-velocity with a contravariant index; it is convenient to consider the covariant four-velocity locally averaged over the particle system as the four-potential of the acceleration field  $u_\mu = \left( \frac{\mathcal{G}}{c}, -\mathbf{U} \right)$ , where  $\mathcal{G}$

and  $\mathbf{U}$  denote the scalar and vector potentials, respectively,

$u_{\mu\nu} = \nabla_\mu u_\nu - \nabla_\nu u_\mu = \partial_\mu u_\nu - \partial_\nu u_\mu$  – the acceleration tensor calculated through the derivatives of the four-potential of the acceleration field,

$\eta$  – a function of coordinates and time,

$\pi_\mu = \frac{p_0}{\rho_0 c^2} u_\mu = \left( \frac{\wp}{c}, -\mathbf{\Pi} \right)$  – the four-potential of the pressure field, consisting of the scalar potential  $\wp$  and the vector potential  $\mathbf{\Pi}$ ,  $p_0$  is the pressure in the reference frame associated with the particle, the relation  $\frac{p_0}{\rho_0 c^2}$  specifies the equation of the matter state,

$f_{\mu\nu} = \nabla_\mu \pi_\nu - \nabla_\nu \pi_\mu = \partial_\mu \pi_\nu - \partial_\nu \pi_\mu$  – the tensor of the pressure field,

$\sigma$  – a function of coordinates and time,

$\sqrt{-g} d\Sigma = \sqrt{-g} c dt dx^1 dx^2 dx^3$  – the invariant four-volume, expressed through the differential of the time coordinate

$dx^0 = c dt$ , through the product  $dx^1 dx^2 dx^3$  of differentials of the spatial coordinates and through the square root  $\sqrt{-g}$  of the determinant  $g$  of the metric tensor, taken with a negative sign.

Action function (1) consists of almost the same terms as those which were considered in [3]. The difference is that now we replace the term with the energy density of particles with four terms located at the end of (1). It is natural to assume that each term is included in (1) relatively independently of the other terms, describing the state of the system in one way or another. The value of the four-potential  $u_\mu$  of the set of matter units or point particles of the system defines the four-field of the system's velocities, and the product  $u_\mu J^\mu$  in (1) can be regarded as the energy of interaction of the mass current  $J^\mu$  with the field of velocities. Similarly,  $D_\mu$  is the four-potential of the gravitational field, and the product  $D_\mu J^\mu$  defines the energy of interaction of the mass current with the gravitational field. The electromagnetic field is specified by the four-potential  $A_\mu$ , the source of the field is the electromagnetic current  $j^\mu$ , and the product of these quantities  $A_\mu j^\mu$  is the density of the energy of interaction of a moving charged matter unit with the electromagnetic field. The invariant of the gravitational field in the form of the tensor product  $\Phi_{\mu\nu} \Phi^{\mu\nu}$  is associated with the gravitational field energy and cannot be equal to zero even outside bodies. The same holds for the electromagnetic field invariant  $F_{\mu\nu} F^{\mu\nu}$ . This follows from the properties of long-range action of the specified fields. As for the field of velocities  $u_\mu$ , the field should be used to describe the motion of the matter particles. Accordingly, the field of accelerations in the form of the tensor  $u_{\mu\nu}$  and the energy of this field associated with the invariant  $u_{\mu\nu} u^{\mu\nu}$  refer to the accelerated motion of particles and are calculated for those spatial points within the system's volume where the matter is located.

The last two terms in (1) are associated with the pressure in the matter, and the product  $\pi_\mu J^\mu$  characterizes the interaction of the pressure field with the mass four-current, and the invariant  $f_{\mu\nu} f^{\mu\nu}$  is part of the stress-energy tensor of the pressure field.

We will also note the difference of four-currents  $J^\mu$  and  $j^\mu$  – all particles of the system make contribution to the mass current  $J^\mu$ , and only charged particles make contribution to the electromagnetic current  $j^\mu$ . This results in difference of the fields' influence – the gravitational field influences any particles and the electromagnetic field influences only the charged particles or the matter, in which by the field can sufficiently divide with its influence the charges of opposite signs from each other. The field of velocities  $u_\mu$ , as well as the mass current  $J^\mu$ , are associated with all the particles of the system. Therefore, the product  $u_\mu J^\mu$  describes that part of the particles' energy, which stays if we somehow "turn off" in the system under consideration all the macroscopic gravitational and electromagnetic fields and remove the pressure, without changing the field of velocities  $u_\mu$  or the mass current  $J^\mu = \rho_0 u^\mu$ .

### Variations of the action function

We will vary the action function  $S$  in (1) term by term, then the total variation  $\delta S$  will be the sum of variations of individual terms. In total, there are 9 terms inside the integral in (1). If we consider the quantity  $\Lambda$  a constant (a cosmological constant), then according to [7-9] the variation of the first term in the action function (1) is equal to:

$$\delta S_1 = \int \left( \begin{array}{c} -k R^{\alpha\beta} + \frac{k}{2} R g^{\alpha\beta} - \\ -k \Lambda g^{\alpha\beta} \end{array} \right) \delta g_{\alpha\beta} \sqrt{-g} d\Sigma, \quad (2)$$

where:

$R^{\alpha\beta}$  is the Ricci tensor,

$\delta g_{\alpha\beta}$  is the variation of the metric tensor.

According to [3], the variations of terms 2 and 3 in the action function are as follows:

$$\delta S_2 = \int \left( \begin{array}{c} -\frac{1}{c} \Phi_{\beta\sigma} J^\sigma \xi^\beta - \frac{1}{c} J^\beta \delta D_\beta \\ -\frac{1}{2c} D_\mu J^\mu g^{\alpha\beta} \delta g_{\alpha\beta} \end{array} \right) \sqrt{-g} d\Sigma, \quad (3)$$

$$\delta S_3 = \int \left( \begin{array}{c} -\frac{c}{4\pi G} \nabla_\alpha \Phi^{\alpha\beta} \delta D_\beta - \\ -\frac{1}{2c} U^{\alpha\beta} \delta g_{\alpha\beta} \end{array} \right) \sqrt{-g} d\Sigma, \quad (4)$$

where:

$\xi^\beta$  is the variation of coordinates, which results in the variation of the mass four-current  $J^\mu$  and in the variation of the electromagnetic four-current  $j^\mu$ ,

$\delta D_\beta$  is the variation of the four-potential of the gravitational field,

and  $U^{\alpha\beta}$  denotes the stress-energy tensor of the gravitational field:

$$\begin{aligned} U^{\alpha\beta} &= \frac{c^2}{4\pi G} \left( g^{\alpha\nu} \Phi_{\kappa\nu} \Phi^{\kappa\beta} - \frac{1}{4} g^{\alpha\beta} \Phi_{\mu\nu} \Phi^{\mu\nu} \right) \\ &= -\frac{c^2}{4\pi G} \left( \Phi^\alpha{}_\kappa \Phi^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} \Phi_{\mu\nu} \Phi^{\mu\nu} \right). \end{aligned} \quad (5)$$

Variations of terms 4 and 5 in the action function according to [6-7] and [10] are as follows:

$$\delta S_4 = \int \left( \begin{array}{c} -\frac{1}{c} F_{\beta\sigma} j^\sigma \xi^\beta - \frac{1}{c} j^\beta \delta A_\beta \\ -\frac{1}{2c} A_\mu j^\mu g^{\alpha\beta} \delta g_{\alpha\beta} \end{array} \right) \sqrt{-g} d\Sigma, \quad (6)$$

$$\delta S_5 = \int \left( \begin{array}{c} c \varepsilon_0 \nabla_\alpha F^{\alpha\beta} \delta A_\beta - \\ -\frac{1}{2c} W^{\alpha\beta} \delta g_{\alpha\beta} \end{array} \right) \sqrt{-g} d\Sigma, \quad (7)$$

where  $\delta A_\beta$  is the variation of the four-potential of the electromagnetic field and  $W^{\alpha\beta}$  denotes

the stress-energy tensor of the electromagnetic field:

$$\begin{aligned} W^{\alpha\beta} &= \varepsilon_0 c^2 \left( -g^{\alpha\nu} F_{\kappa\nu} F^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} F_{\mu\nu} F^{\mu\nu} \right) \\ &= \varepsilon_0 c^2 \left( F^\alpha_\kappa F^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} F_{\mu\nu} F^{\mu\nu} \right). \end{aligned} \quad (8)$$

Variations of the other terms in the action function (1) are defined in Appendices A-D and have the following form:

$$\delta S_6 = \int \left( \begin{aligned} &-\frac{1}{c} u_{\beta\sigma} J^\sigma \xi^\beta - \frac{1}{c} J^\beta \delta u_\beta \\ &-\frac{1}{2c} u_\mu J^\mu g^{\alpha\beta} \delta g_{\alpha\beta} \end{aligned} \right) \sqrt{-g} d\Sigma, \quad (9)$$

$$\delta S_7 = \int \left( \begin{aligned} &\frac{c}{4\pi\eta} \nabla_\alpha u^{\alpha\beta} \delta u_\beta - \\ &-\frac{1}{2c} B^{\alpha\beta} \delta g_{\alpha\beta} \end{aligned} \right) \sqrt{-g} d\Sigma, \quad (10)$$

where  $\delta u_\beta$  is the variation of the four-potential of the acceleration field and  $B^{\alpha\beta}$  denotes the stress-energy tensor of the field of accelerations:

$$B^{\alpha\beta} = \frac{c^2}{4\pi\eta} \left( -g^{\alpha\nu} u_{\kappa\nu} u^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} u_{\mu\nu} u^{\mu\nu} \right). \quad (11)$$

$$\delta S_8 = \int \left( \begin{aligned} &-\frac{1}{c} f_{\beta\sigma} J^\sigma \xi^\beta - \frac{1}{c} J^\beta \delta \pi_\beta \\ &-\frac{1}{2c} \pi_\mu J^\mu g^{\alpha\beta} \delta g_{\alpha\beta} \end{aligned} \right) \sqrt{-g} d\Sigma. \quad (12)$$

$$\delta S_9 = \int \left( \begin{aligned} &\frac{c}{4\pi\sigma} \nabla_\alpha f^{\alpha\beta} \delta \pi_\beta - \\ &-\frac{1}{2c} P^{\alpha\beta} \delta g_{\alpha\beta} \end{aligned} \right) \sqrt{-g} d\Sigma, \quad (13)$$

where the stress-energy tensor of the pressure field is given as:

$$P^{\alpha\beta} = \frac{c^2}{4\pi\sigma} \left( -g^{\alpha\nu} f_{\kappa\nu} f^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} f_{\mu\nu} f^{\mu\nu} \right). \quad (14)$$

In variation, in order to simplify (10), the special case is considered, when  $\eta$  is a constant, which does not vary by definition. According to its meaning,  $\eta$  depends on the parameters of the system under consideration and can therefore have different values. The same should be said about  $\sigma$ .

## The Motion Equations of the Field, Particles and Metric

According to the principle of least action, we should sum up all the variations of the individual terms of the action function and equate the result to zero. The sum of variations (2), (3), (4), (6), (7), (9), (10), (12) and (13) gives the total variation of the action function:

$$\begin{aligned} \delta S &= \delta S_1 + \delta S_2 + \delta S_3 + \delta S_4 + \delta S_5 \\ &+ \delta S_6 + \delta S_7 + \delta S_8 + \delta S_9 = 0. \end{aligned} \quad (15)$$

### The field equations

When the system moves in spacetime, the variations  $\delta g_{\alpha\beta}$ ,  $\xi^\beta$ ,  $\delta D_\beta$ ,  $\delta A_\beta$ ,  $\delta u_\beta$  and  $\delta \pi_\beta$  do not vanish, since it is supposed that they can occur only at the beginning and the end of the process, when the conditions of motion are precisely fixed. Consequently, the sum of the terms, which is located before these variations, should vanish. For example, the variation  $\delta A_\beta$  occurs only in  $\delta S_4$  according to (6) and in  $\delta S_5$  from (7), then from (15) it follows that:

$$\int \left( -\frac{1}{c} j^\beta + c \varepsilon_0 \nabla_\alpha F^{\alpha\beta} \right) \delta A_\beta \sqrt{-g} d\Sigma = 0.$$

From this, we obtain the equation of the electromagnetic field with the field sources as:

$$\nabla_\alpha F^{\alpha\beta} = \frac{1}{c^2 \varepsilon_0} j^\beta \quad \text{or} \quad \nabla_\beta F^{\alpha\beta} = -\mu_0 j^\alpha, \quad (16)$$

where  $\mu_0 = \frac{1}{c^2 \varepsilon_0}$  is the vacuum permeability.

The second equation of the electromagnetic field follows from the definition of the electromagnetic tensor in terms of the electromagnetic four-potential and from the antisymmetry properties of this tensor:

$$\nabla_\sigma F_{\mu\nu} + \nabla_\nu F_{\sigma\mu} + \nabla_\mu F_{\nu\sigma} = 0$$

or  $\varepsilon^{\alpha\beta\gamma\delta} \nabla_\gamma F_{\alpha\beta} = 0$ , (17)

where  $\varepsilon^{\alpha\beta\gamma\delta}$  is a Levi-Civita symbol or a completely antisymmetric unit tensor.

The variation  $\delta D_\beta$  is present only in (3) and (4), so that according to (15) we obtain:

$$\int \left( -\frac{1}{c} J^\beta - \frac{c}{4\pi G} \nabla_\alpha \Phi^{\alpha\beta} \right) \delta D_\beta \sqrt{-g} d\Sigma = 0.$$

The equation of gravitational field with the field sources follows from this as:

$$\nabla_\alpha \Phi^{\alpha\beta} = -\frac{4\pi G}{c^2} J^\beta \text{ or } \nabla_\beta \Phi^{\alpha\beta} = \frac{4\pi G}{c^2} J^\alpha. \quad (18)$$

If we take into account the definition of the gravitational tensor:  $\Phi_{\mu\nu} = \nabla_\mu D_\nu - \nabla_\nu D_\mu = \partial_\mu D_\nu - \partial_\nu D_\mu$  and take the covariant derivative of this tensor with subsequent cyclic interchange of the indices, the following equations are solved identically:

$$\nabla_\sigma \Phi_{\mu\nu} + \nabla_\nu \Phi_{\sigma\mu} + \nabla_\mu \Phi_{\nu\sigma} = 0$$

or  $\varepsilon^{\alpha\beta\gamma\delta} \nabla_\gamma \Phi_{\alpha\beta} = 0$ . (19)

Equation (19) without the sources and equation (18) with the sources define a complete set of gravitational field equations in the covariant theory of gravitation.

Consider now the rule for the difference of the second covariant derivatives with respect to the covariant derivative of the electromagnetic four-potential  $\nabla^\alpha A^\beta$ :

$$\begin{aligned} (\nabla_\alpha \nabla_\beta - \nabla_\beta \nabla_\alpha) \nabla^\alpha A^\beta &= \\ -R_{\mu,\alpha\beta}^\alpha \nabla^\mu A^\beta - R_{\mu,\alpha\beta}^\beta \nabla^\alpha A^\mu &= \\ = R_{\mu,\alpha\beta}^\beta (\nabla^\mu A^\alpha - \nabla^\alpha A^\mu) &= -R_{\mu\alpha} F^{\mu\alpha}. \end{aligned}$$

With the rule in mind, the application of the covariant derivative  $\nabla_\alpha$  to (16) and (18) gives the following:

$$\begin{aligned} \nabla_\alpha \nabla_\beta F^{\alpha\beta} &= \nabla_\alpha \nabla_\beta \nabla^\alpha A^\beta - \nabla_\alpha \nabla_\beta \nabla^\beta A^\alpha \\ &= -R_{\mu\alpha} F^{\mu\alpha} = -\mu_0 \nabla_\alpha j^\alpha. \end{aligned}$$

$$R_{\mu\alpha} \Phi^{\mu\alpha} = -\frac{4\pi G}{c^2} \nabla_\alpha J^\alpha.$$

This shows that field tensors  $F^{\mu\alpha}$  and  $\Phi^{\mu\alpha}$  lead to the divergence of the corresponding four-currents in a curved space-time. Mixed curvature tensor  $R_{\mu,\alpha\beta}^\beta$  and Ricci tensor  $R_{\mu\alpha}$  vanish only in Minkowski space. In this case, the covariant derivatives become the partial derivatives and the continuity equations for the gravitational and electromagnetic four-currents in the special theory of relativity are obtained:

$$\partial_\alpha j^\alpha = 0, \quad \partial_\alpha J^\alpha = 0. \quad (20)$$

We will note that in order to simplify the equations for the four-potential of fields, we can use expressions which are called gauge conditions:

$$\nabla_\beta D^\beta = \nabla^\mu D_\mu = 0, \quad \nabla_\beta A^\beta = \nabla^\mu A_\mu = 0. \quad (21)$$

### The acceleration field equations

The variation of four-potential  $\delta u_\beta$  is included in (9) and (10). Therefore, according to (15), we obtain:

$$\begin{aligned} \int \left( -\frac{1}{c} J^\beta + \frac{c}{4\pi\eta} \nabla_\alpha u^{\alpha\beta} \right) \delta u_\beta \sqrt{-g} d\Sigma &= 0. \\ \nabla_\alpha u^{\alpha\beta} &= \frac{4\pi\eta}{c^2} J^\beta, \text{ or } \nabla_\beta u^{\alpha\beta} = -\frac{4\pi\eta}{c^2} J^\alpha. \end{aligned} \quad (22)$$

If we compare (18) and (22), it turns out that the presence of the four-vector of mass current  $J^\alpha$  not only leads to the occurrence of space-time gradient of the gravitational field in the system under consideration, but also is generally accompanied by changes in time or by four-velocity gradients of the particles that constitute this system. Besides, the covariant four-velocities of the whole set of particles forms the velocity field  $u_\beta$ , the derivatives of which define the acceleration field and are described by the tensor  $u_{\mu\nu}$ . As an example of a system, where it can be clearly observed, we can take a rotating partially-charged collapsing gas-dust cloud, held by gravity. An ordered acceleration field occurs in the cloud due to the rotational acceleration and contains the centripetal and tangential acceleration.

Due to its definition in the form of a four-rotor of  $u_\beta$ , the following relations hold for acceleration tensor  $u_{\mu\nu}$ :

$$\begin{aligned} \nabla_\sigma u_{\mu\nu} + \nabla_\nu u_{\sigma\mu} + \nabla_\mu u_{\nu\sigma} &= 0 \\ \text{or } \varepsilon^{\alpha\beta\gamma\delta} \nabla_\gamma u_{\alpha\beta} &= 0. \end{aligned} \quad (23)$$

As we can see, the structure of equations (22) and (23) for the acceleration field is similar to the structure of equations for the strengths of gravitational and electromagnetic fields.

In the local geodetic reference frame, the derivatives of the metric tensor and the curvature tensor become equal to zero, the covariant derivative becomes a partial derivative and the equations take the simplest form. We will go over to this reference frame and apply the derivative  $\partial^\nu$  to (23) and make substitution for the first and third terms using (22):

$$\begin{aligned} 0 &= \partial_\sigma \partial^\nu u_{\mu\nu} + \partial^\nu \partial_\nu u_{\sigma\mu} + \partial_\mu \partial^\nu u_{\nu\sigma} \\ &= \partial_\sigma \left( -\frac{4\pi\eta}{c^2} J_\mu \right) + \square u_{\sigma\mu} + \partial_\mu \left( \frac{4\pi\eta}{c^2} J_\sigma \right). \\ \square u_{\sigma\mu} &= \partial_\sigma \left( \frac{4\pi\eta}{c^2} J_\mu \right) - \partial_\mu \left( \frac{4\pi\eta}{c^2} J_\sigma \right). \end{aligned}$$

If we apply definition  $u_{\sigma\mu} = \partial_\sigma u_\mu - \partial_\mu u_\sigma$  to four-d'Alembertian  $\square u_{\sigma\mu}$ , where  $\square = \partial^\nu \partial_\nu$  is the d'Alembert operator, it will give:

$$\square u_{\sigma\mu} = \square (\partial_\sigma u_\mu - \partial_\mu u_\sigma) = \partial_\sigma \square u_\mu - \partial_\mu \square u_\sigma.$$

Comparing with the previous expression, we find the wave equation for the four-potential  $u_\mu$ :

$$\square u_\mu = \frac{4\pi\eta}{c^2} J_\mu.$$

On the other hand, after lowering the acceleration tensor indices, we have from (22):

$$\begin{aligned} \frac{4\pi\eta}{c^2} J_\mu &= \partial^\nu u_{\nu\mu} = \partial^\nu (\partial_\nu u_\mu - \partial_\mu u_\nu) \\ &= \square u_\mu - \partial_\mu \partial^\nu u_\nu. \end{aligned}$$

Comparing this equation with equation for  $\square u_\mu$  leads to the expression:

$$\partial^\nu u_\nu = \partial_\mu u^\mu = \varsigma,$$

where  $\varsigma$  is some constant.

In an arbitrary reference frame, we should specify the obtained expressions, since in contrast to permutations of partial derivatives, in case of permutation of the covariant derivatives from the sequence  $\nabla_\mu \nabla_\nu$  to the sequence  $\nabla_\nu \nabla_\mu$ , some additional terms appear. In particular, if we use the relation:

$$\nabla^\nu u_\nu = \nabla_\mu u^\mu = 0, \quad (24)$$

then after substituting the expression  $u^{\alpha\beta} = \nabla^\alpha u^\beta - \nabla^\beta u^\alpha$  in (22), the wave equation can be written as follows:

$$\left. \begin{aligned} &g^{\rho\nu} \partial_\rho \partial_\nu u^\alpha \\ &+ g^{\rho\nu} \left( \Gamma_{\nu s}^\alpha \partial_\rho u^s - \Gamma_{\rho\nu}^s \partial_s u^\alpha \right. \\ &\quad \left. + \Gamma_{s\rho}^\alpha \partial_\nu u^s + u^s \partial_s \Gamma_{\nu\rho}^\alpha \right) \\ &= \frac{4\pi\eta}{c^2} J^\alpha. \end{aligned} \right\} \quad (25)$$

In the curved space, operator  $\nabla_\nu \nabla^\nu$  acts differently on scalars, four-vectors and four-tensors, and it usually contains the Ricci tensor. Due to condition (24), the Ricci tensor is absent in (25), but the terms with the Christoffel symbols remain.

Equation (24) is a gauge condition for the four-potential  $u_\nu$ , which is similar by its meaning to gauge conditions (21) for the electromagnetic and gravitational four-potentials. Both (24) and (25) will hold on condition that  $\eta = \text{const}$ .

In Appendix E, it will be shown that the acceleration tensor  $u_{\mu\nu}$  includes the vector components **S** and **N**, based on which, according to (E6), we can build a four-vector of particles' acceleration.

### The pressure field equations

To obtain the pressure field equations, we need to choose in (15) those terms which contain the variation  $\delta\pi_\beta$ . This variation is present in (12) and (13), which gives the following:

$$\int \left( -\frac{1}{c} J^\beta + \frac{c}{4\pi\sigma} \nabla_\alpha f^{\alpha\beta} \right) \delta\pi_\beta \sqrt{-g} d\Sigma = 0.$$

$$\nabla_\alpha f^{\alpha\beta} = \frac{4\pi\sigma}{c^2} J^\beta \quad \text{or} \quad \nabla_\beta f^{\alpha\beta} = -\frac{4\pi\sigma}{c^2} J^\alpha. \quad (26)$$

It follows from (26) that the mass four-current generates the pressure field in bodies, which can be described by the pressure tensor  $f^{\alpha\beta}$ . The same relations hold for this tensor as for the tensors of other fields:

$$\begin{aligned} \nabla_\sigma f_{\mu\nu} + \nabla_\nu f_{\sigma\mu} + \nabla_\mu f_{\nu\sigma} &= 0 \\ \text{or } \varepsilon^{\alpha\beta\gamma\delta} \nabla_\gamma f_{\alpha\beta} &= 0. \end{aligned} \quad (27)$$

The wave equation for the four-potential of the pressure field follows from (26):

$$\left. \begin{aligned} &g^{\rho\nu} \partial_\rho \partial_\nu \pi^\alpha + \\ &g^{\rho\nu} \left( \Gamma_{\nu s}^\alpha \partial_\rho \pi^s - \Gamma_{\rho\nu}^s \partial_s \pi^\alpha \right. \\ &\quad \left. + \Gamma_{s\rho}^\alpha \partial_\nu \pi^s + \pi^s \partial_s \Gamma_{\nu\rho}^\alpha \right) \\ &= \frac{4\pi\sigma}{c^2} J^\alpha. \end{aligned} \right\} \quad (28)$$

Equation (28) will be valid if there is gauge condition of the pressure four-potential:

$$\nabla^\nu \pi_\nu = \nabla_\mu \pi^\mu = 0. \quad (29)$$

The properties of the pressure field are described in Appendix F, where it is shown that the pressure tensor  $f_{\mu\nu}$  contains two vector components **C** and **I**, which determine the energy and the pressure force, as well as the pressure energy flux.

### The equations of motion of particles

The variation  $\xi^\beta$  that leads to the equations of motion of the particles is present in (3), (6), (9) and (12). For this variation, it follows from (15):

$$\begin{aligned} &\int \left( -\frac{1}{c} \Phi_{\beta\sigma} J^\sigma - \frac{1}{c} F_{\beta\sigma} j^\sigma - \right. \\ &\quad \left. -\frac{1}{c} u_{\beta\sigma} J^\sigma - \frac{1}{c} f_{\beta\sigma} J^\sigma \right) \xi^\beta \sqrt{-g} d\Sigma = 0, \\ &-u_{\beta\sigma} J^\sigma = \Phi_{\beta\sigma} J^\sigma + F_{\beta\sigma} j^\sigma + f_{\beta\sigma} J^\sigma. \end{aligned}$$

The left side of the equation can be transformed, considering the expression  $J^\sigma = \rho_0 u^\sigma$  for the four-vector of mass current

density and the definition of the acceleration tensor  $u_{\beta\sigma} = \nabla_\beta u_\sigma - \nabla_\sigma u_\beta$ :

$$\left. \begin{aligned} -u_{\beta\sigma} J^\sigma &= -\rho_0 u^\sigma (\nabla_\beta u_\sigma - \nabla_\sigma u_\beta) \\ &= \rho_0 u^\sigma \nabla_\sigma u_\beta = \rho_0 \frac{D u_\beta}{D\tau}. \end{aligned} \right\} \quad (30)$$

We used the relation  $u^\sigma \nabla_\beta u_\sigma = 0$ , which follows from the equation  $\nabla_\beta (u^\sigma u_\sigma) = \nabla_\beta (c^2) = 0$  and the operator of proper-time-derivative as operator of the derivative with respect to the proper time  $u^\sigma \nabla_\sigma = \frac{D}{D\tau}$ , where  $D$  is a symbol of four-

differential in curved spacetime and  $\tau$  is the proper time [11]. Taking (30) into account, the equation of motion takes the form:

$$\rho_0 \frac{D u_\beta}{D\tau} = \Phi_{\beta\sigma} J^\sigma + F_{\beta\sigma} j^\sigma + f_{\beta\sigma} J^\sigma. \quad (31)$$

We will note that the equations of field motion (16) – (19), of the acceleration field (22) and (23), of the pressure field (26) and (27) and the equation of the particles' motion (31) are differential equations, which are valid at any point volume of spacetime in the system under consideration. In particular, if the mass density  $\rho_0$  in some point volume is zero, then all the terms in (31) will be zero.

The quantity  $\frac{D u_\beta}{D\tau}$  at the left side of (31) is

the four-acceleration of a point particle, while the proper time differential  $D\tau = d\tau$  is associated with the interval by relation:  $ds = c d\tau$  and the relation:  $D u_\beta = dx^\sigma \nabla_\sigma u_\beta$  holds. The first two terms at the right side of (31) are the densities of the gravitational and electromagnetic four-forces, respectively. It can be shown (see for example [3], [12]) that for four-forces exerted by the field on the particle, there are alternative expressions in terms of the stress-energy tensors (5) and (8):

$$\Phi_{\beta\sigma} J^\sigma = -\nabla^k U_{\beta k}, \quad F_{\beta\sigma} j^\sigma = -\nabla^k W_{\beta k}. \quad (32)$$

Similarly, the left side of (31) with regard to (30) is expressed in terms of stress-energy tensor of the acceleration field (11) as:



$$\rho_0 \frac{Du_\beta}{D\tau} = -u_{\beta\sigma} J^\sigma = \nabla^k B_{\beta k}. \quad (33)$$

To prove (33), we should expand the tensor  $B_{\beta k}$  with the help of definition (11), apply the covariant derivative  $\nabla^k$  to the tensor products and then use equations (22) and (23). Equation (33) shows that the four-acceleration of the particle can be described by either the acceleration tensor  $u_{\beta\sigma}$  or the tensor  $B_{\beta k}$ .

For the pressure field, we can write the same as for other fields:

$$f_{\beta\sigma} J^\sigma = -\nabla^k P_{\beta k}. \quad (34)$$

In (34), the pressure four-force is associated with the covariant derivative of the stress-energy tensor of the pressure field.

From (31) – (34) it follows that:

$$\begin{aligned} \nabla^k (B_{\beta k} + U_{\beta k} + W_{\beta k} + P_{\beta k}) &= 0 \\ \text{or } \nabla_\beta (B^{\alpha\beta} + U^{\alpha\beta} + W^{\alpha\beta} + P^{\alpha\beta}) &= 0. \end{aligned} \quad (35)$$

In Minkowski space  $u_\beta = (\gamma c, -\gamma \mathbf{v})$ , where

$$\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$$

is present, four-differentials  $D$  become ordinary differentials  $d$ ,  $D\tau = d\tau = dt/\gamma$ , and the motion equation (31) falls into the scalar and vector equations, while the vector equation contains the total gravitational force with regard to the torsion field, the electromagnetic Lorentz force and the pressure force:

$$\rho_0 c^2 \frac{d\gamma}{dt} = \rho_0 \mathbf{v} \cdot \mathbf{\Gamma} + \rho_{0q} \mathbf{v} \cdot \mathbf{E} + \rho_0 \mathbf{v} \cdot \mathbf{C}, \quad (36)$$

$$\begin{aligned} & \int \left( -kR^{\alpha\beta} + \frac{k}{2} R g^{\alpha\beta} - k\Lambda g^{\alpha\beta} - \frac{1}{2c} D_\mu J^\mu g^{\alpha\beta} - \frac{1}{2c} U^{\alpha\beta} - \frac{1}{2c} A_\mu j^\mu g^{\alpha\beta} \right. \\ & \quad \left. - \frac{1}{2c} W^{\alpha\beta} - \frac{1}{2c} u_\mu J^\mu g^{\alpha\beta} - \frac{1}{2c} B^{\alpha\beta} - \frac{1}{2c} \pi_\mu J^\mu g^{\alpha\beta} - \frac{1}{2c} P^{\alpha\beta} \right) \delta g_{\alpha\beta} \sqrt{-g} d\Sigma = 0. \\ & -2ckR^{\alpha\beta} + ckR g^{\alpha\beta} - 2ck\Lambda g^{\alpha\beta} = \\ & = D_\mu J^\mu g^{\alpha\beta} + U^{\alpha\beta} + A_\mu j^\mu g^{\alpha\beta} + W^{\alpha\beta} + u_\mu J^\mu g^{\alpha\beta} + B^{\alpha\beta} + \pi_\mu J^\mu g^{\alpha\beta} + P^{\alpha\beta}. \end{aligned} \quad (38)$$

The equation for the metric (38) allows to determine the metric tensor  $g^{\alpha\beta}$  by the known quantities characterizing the matter and field. If

$$\left. \begin{aligned} \rho_0 \frac{d(\gamma \mathbf{v})}{dt} &= \rho_0 (\mathbf{\Gamma} + [\mathbf{v} \times \mathbf{\Omega}]) \\ &+ \rho_{0q} (\mathbf{E} + [\mathbf{v} \times \mathbf{B}]) + \rho_0 (\mathbf{C} + [\mathbf{v} \times \mathbf{I}]), \end{aligned} \right\} \quad (37)$$

where  $\mathbf{v}$  is the velocity of a point particle,  $\mathbf{\Gamma}$  is the gravitational field strength,  $\rho_{0q}$  is the charge density,  $\mathbf{E}$  is the electric field strength,  $\mathbf{C}$  is the pressure field strength,  $\mathbf{\Omega}$  is the torsion field vector,  $\mathbf{B}$  is the magnetic field induction and  $\mathbf{I}$  is the solenoidal vector of the pressure field.

If during the time  $dt$  the density  $\rho_0$  does not change, it can be put under the derivative's sign. Then, at the left side of (36) the quantity

$$\frac{dE_r}{dt} \text{ appears, where } E_r = \frac{\rho_0 c^2}{\sqrt{1-v^2/c^2}}$$

is the relativistic energy density. Similarly, at the left

side of (37) the quantity  $\frac{d\mathbf{J}}{dt}$  appears, where

$$\mathbf{J} = \frac{\rho_0 \mathbf{v}}{\sqrt{1-v^2/c^2}} \text{ is the mass three-current density.}$$

### The equations for the metric

Let us consider action variations (2), (3), (4), (6), (7), (9), (10), (12) and (13), which contain the variation  $\delta g_{\alpha\beta}$ . The sum of all the terms in (15) with the variation  $\delta g_{\alpha\beta}$  must be zero:

we take the covariant derivative  $\nabla_\beta$  in this equation, the left side of the equation vanishes

on condition  $\Lambda = \text{const}$ . Taking (35) into account, we obtain:

$$\nabla_\beta \left( D_\mu J^\mu g^{\alpha\beta} + A_\mu j^\mu g^{\alpha\beta} + u_\mu J^\mu g^{\alpha\beta} + \pi_\mu J^\mu g^{\alpha\beta} \right) = 0, \quad (39)$$

where  $\chi$  is a function of time and coordinates and the scalar invariant with respect to coordinate transformations.

If we expand the scalar products of vectors using the expressions:

$$\begin{aligned} D_\mu J^\mu &= c \rho_0 \frac{dt}{ds} D_\mu \frac{dx^\mu}{dt} = c \rho_0 \frac{dt}{ds} (\psi - \mathbf{v} \cdot \mathbf{D}), \\ A_\mu j^\mu &= c \rho_{0q} \frac{dt}{ds} (\varphi - \mathbf{v} \cdot \mathbf{A}), \\ \pi_\mu J^\mu &= c \rho_0 \frac{dt}{ds} (\wp - \mathbf{v} \cdot \mathbf{\Pi}), \end{aligned} \quad (40)$$

then (39) can be written as:

$$\left. \begin{aligned} c \rho_0 \frac{dt}{ds} (\psi - \mathbf{v} \cdot \mathbf{D}) + c \rho_{0q} \frac{dt}{ds} (\varphi - \mathbf{v} \cdot \mathbf{A}) \\ + \rho_0 c^2 + c \rho_0 \frac{dt}{ds} (\wp - \mathbf{v} \cdot \mathbf{\Pi}) = \chi. \end{aligned} \right\} \quad (41)$$

If the system's matter and charges are divided into small pieces and scattered to infinity, then there the external field potentials become equal to zero, since interparticle interaction tends to be zero, and at  $\mathbf{v} = 0$  we obtain the following:

$$(\rho_0 \psi_0 + \rho_{0q} \varphi_0 + \rho_0 c^2 + p_0)_\infty = \chi. \quad (42)$$

Consequently,  $\chi$  is associated with the particle's proper scalar potentials  $\psi_0$  and  $\varphi_0$ , the mass density  $\rho_0$  and the pressure  $p_0$  in the particle located at infinity. Expression (41) can be considered as the differential law of conservation of mass-energy: the greater the velocity  $\mathbf{v}$  of a point particle is, and the greater the gravitational field potentials  $\psi$  and  $\mathbf{D}$ , the electromagnetic field potentials  $\varphi$  and  $\mathbf{A}$  and the pressure field potentials  $\wp$  and  $\mathbf{\Pi}$  are, the more the mass density  $\rho_0$  differs from its value at infinity. For example, if a point particle falls into the gravitational field with the potential  $\psi$ , then the change in the particle's energy is

described by the term  $c \frac{dt}{ds} \rho_0 \psi$ . According to (41), such energy change can be compensated by the change in the rest energy of the particle due to the change  $\rho_0$ . Since the gravitational field potential  $\psi$  is always negative, then the mass density  $\rho_0$  and the pressure inside the point particle should increase due to the field potential.

This is possible, if we remember that the whole procedure of deriving the motion equations of particles, field and metric from the principle of least action is based on the fact that the mass and charge of the matter unit at varying of the coordinates remain constant, despite of the change in the charge density, mass density and its volume [7]. If the mass of a simple system in the form of a point particle and the fields associated with it is proportional to  $\chi$ , then according to (41) the mass of such a system remains unchanged, despite of the change in the fields, mass density  $\rho_0$  and pressure  $p_0$ . Conservation of the mass-energy of each particle with regard to the mass-energy of the fields leads to conservation of the mass-energy of an arbitrary system including a multitude of particles and the fields surrounding them. We will remind that this article refers to the continuously distributed matter, so that each point particle or a unit of this matter may have its own mass density  $\rho_0$  and its value  $\chi$ .

We will now return to (38) and take the contraction of tensors by means of multiplying the equation by  $g_{\alpha\beta}$ , taking into account the relation  $g_{\alpha\beta} g^{\alpha\beta} = 4$ , and then dividing all by 2:

$$c k R - 4 c k \Lambda = \left. \begin{aligned} 2 D_\mu J^\mu + 2 A_\mu j^\mu + 2 u_\mu J^\mu + 2 \pi_\mu J^\mu \end{aligned} \right\}, \quad (43)$$

where  $R = g_{\alpha\beta} R^{\alpha\beta}$  is the scalar curvature, and it was taken into account that the contractions of tensors  $U^{\alpha\beta}$ ,  $W^{\alpha\beta}$ ,  $B^{\alpha\beta}$  and  $P^{\alpha\beta}$  are equal to zero.

In case if the cosmological constant  $\Lambda$  were known, based on (43), we could find the scalar curvature  $R$ .

In order to simplify equation (38) in [3] and [4], we introduced the gauge for  $\Lambda$ , at which the following equation would hold, if we additionally take into account the term with the pressure  $\pi_\mu J^\mu$ :

$$2ck\Lambda = -u_\mu J^\mu - D_\mu J^\mu - A_\mu j^\mu - \pi_\mu J^\mu. \quad (44)$$

In the gauge (44), the equation for the metric (38) takes the following form, provided that  $-2ck = \frac{c^4}{8\pi G\beta}$ , where  $\beta$  is a constant of order of unity:

$$\left. \begin{aligned} R^{\alpha\beta} - \frac{1}{2}R g^{\alpha\beta} = \\ \frac{8\pi G\beta}{c^4} (U^{\alpha\beta} + W^{\alpha\beta} + B^{\alpha\beta} + P^{\alpha\beta}). \end{aligned} \right\} \quad (45)$$

We will note that if from the right side of (45) we exclude the stress-energy tensor of the gravitational field  $U^{\alpha\beta}$ , replace the tensor  $B^{\alpha\beta}$  by the stress-energy tensor of the matter in the form  $\phi^{\alpha\beta} = \rho_0 u^\alpha u^\beta$  and neglect the tensor  $P^{\alpha\beta}$ , then at  $\beta=1$  we will obtain a typical equation for the metric used in the general theory of relativity:

$$R^{\alpha\beta} - \frac{1}{2}R g^{\alpha\beta} = \frac{8\pi G}{c^4} (W^{\alpha\beta} + \phi^{\alpha\beta}). \quad (46)$$

The equation for the metric (38) and expression (39) must hold in the covariant theory of gravitation, provided that  $\Lambda = \text{const}$ . If in (39) we remove the term  $D_\mu J^\mu$ , then we will obtain an expression suitable for use in the general theory of relativity. In this case, given that  $u_\mu J^\mu = \rho_0 c^2$ , instead of (39), we obtain the following:

$$A_\mu j^\mu + \rho_0 c^2 + \pi_\mu J^\mu = \chi. \quad (47)$$

If in (47) we equate the term with the energy of particles in the electromagnetic field (in the case when the field is zero) to zero, then the sum of the rest energy density and the pressure energy of each uncharged point particle must be unchanged. It follows that the pressure change must be accompanied by a change in the mass density. If the system contains the electromagnetic field with the four-potential  $A_\mu$  acting on the four-currents  $j^\mu$  generating them,


then in the general case there must be an inverse correlation of the rest energy, pressure energy and the energy of charges in the electromagnetic field.

Indeed, in the general theory of relativity, the mass density determines the rest energy density and spacetime metric, which represents the gravitational field. In (47), the energy of charges in the electromagnetic field is specified by the term  $A_\mu j^\mu$ , and the mass density and hence the metric are associated with this energy at a constant  $\chi$ . On the other hand, the metric is obtained from (46). Therefore, the occurrence of the electromagnetic field influences the metric in two relations — in (47) the mass density and the corresponding metric change, as well as in the equation for the metric (46) the stress-energy tensor of the matter  $\phi^{\alpha\beta}$  changes, while the stress-energy tensor of the electromagnetic field  $W^{\alpha\beta}$  also makes contribution to the metric.

A well-known paradox of general theory of relativity is associated with all of this — the electromagnetic field influences the density, the mass of the bodies as the source of gravitation and the metric, while the gravitational field itself (i.e., metric) does not influence the electrical charges of the bodies, which are the sources of the electromagnetic field. Thus the gravitational and electromagnetic fields are unequal relative to each other, despite the similarity of field equations and the same character of long-range action. Above, we pointed out the fact that the mass four-current leads to the gravitational field gradients, and the addition of the charge to this mass current generates additional electromagnetic (charge) four-current and the corresponding electromagnetic field gradients, depending on the sign of the charge. From this, we can see that the gravitational field looks like a fundamental, basic and indestructible field and the electromagnetic field manifests as some superstructure and the result of charge separation in the initially neutral matter.

If we consider (44) to be valid, then from comparison with (39), we see that the equation  $2ck\Lambda = -\chi$  must be satisfied. Thus, when  $\Lambda$  is considered as a cosmological constant, we can use it to achieve simplification of the equation for the metric (38) and bring it to the form of (45). At the same time, the relation (39) is symmetrical with respect to the contribution of the gravitational and electromagnetic fields to

the density, in spite of the difference in fields. We will remind that in the equation of motion (31), both fields also make symmetrical contributions to the four-acceleration of a point charge.

Although the gauge for  $\Lambda$  in the form of (44) seems the simplest and simplifies some of the equations, in Section 7 the necessity and convenience of another gauge will be shown. 

## Hamiltonian

In this and the next sections, we rely on the standard approach of analytical mechanics. As the coordinates, it is convenient to choose a set of Cartesian coordinates:  $x^0 = ct$ ,  $x^1 = x$ ,  $x^2 = y$ ,  $x^3 = z$ .

Let us consider action (1) and express the Lagrangian from it:

$$L = \int \left( -u_\mu J^\mu - D_\mu J^\mu - A_\mu j^\mu - \pi_\mu J^\mu \right) \sqrt{-g} dx^1 dx^2 dx^3 + \int \left( c k R - 2c k \Lambda + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} dx^1 dx^2 dx^3. \quad (48)$$

The integration in (48) is carried out over the infinite three-dimensional volume of space and over all the material particles of the system. We assume that the scalar curvature  $R$  depends on the metric tensor, and the metric tensor  $g_{\mu\nu}$ , the field tensors  $\Phi_{\mu\nu}$ ,  $F_{\mu\nu}$ ,  $u_{\mu\nu}$ ,  $f_{\mu\nu}$ , the mass density  $\rho_0$ , the charge density  $\rho_{0q}$  and the pressure  $p_0$  are functions of the coordinates  $t, x, y, z$  and do not depend on the particle velocities. Then, the Lagrangian in its general form (48) depends on the coordinates, as well as on the four-potential of pressure  $\pi_\mu$  and four-potentials of the gravitational and electromagnetic fields  $D_\mu$  and  $A_\mu$ .

We will divide the first integral in the Lagrangian (48) into the sum of particular integrals, each of which describes the state of one of the set  $N_p$  of the system's particles. We will also take into account that the Lagrangian depends on the three-dimensional velocities of the particles  $\mathbf{v} = \left( \frac{dx}{dt}, \frac{dy}{dt}, \frac{dz}{dt} \right) = (\dot{x}, \dot{y}, \dot{z})$ , where  $n=1, 2, 3, \dots, N_p$  specifies the particle's number, while the velocity of any particle is part of only one corresponding particular integral. If we denote by  $L_f$  the second integral in (48), which is associated with the energies of fields

inside and outside the fixed physical system and is independent of the particles' velocities, then we can write for the Lagrangian:

$$L = L(t, x, y, z, \mathbf{v}, D_\mu, A_\mu, \pi_\mu) = \sum_{n=1}^{N_p} L^n + L_f = L_n + L_f,$$

where  $L^n = \int \left( -u_\mu J^\mu - D_\mu J^\mu - A_\mu j^\mu - \pi_\mu J^\mu \right) \sqrt{-g} dx^1 dx^2 dx^3$  is a particular Lagrangian of an arbitrary particle.

We will introduce now the Hamiltonian  $H$  of the system as a function of generalized three-dimensional momenta  $\mathbf{P}$  of the particles:

$H = H(t, x, y, z, \mathbf{P}, D_\mu, A_\mu, \pi_\mu)$ . Under the system's generalized momentum, we mean the sum of the generalized momenta of the whole set of particles:

$$\mathbf{P} = \sum_{n=1}^{N_p} \mathbf{P}^n = \mathbf{P}_n = \mathbf{P}^1 + \mathbf{P}^2 + \mathbf{P}^3 + \dots + \mathbf{P}^{N_p}.$$

To find the Hamiltonian, we will apply the Legendre transformations to the system of particles:

$$H = \sum_{n=1}^{N_p} \left( \mathbf{P}^n \cdot \mathbf{v} \right) - L = \left( \mathbf{P} \cdot \mathbf{v} \right) - L, \quad (49)$$

provided that:

$$\mathbf{P} = \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{v}} = \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{v}}. \quad (50)$$

The equality in (50) gives the definition of the generalized momentum  $\mathbf{P}$ , and we can see that the generalized momentum of an arbitrary

particle equals  $\mathbf{P} = \frac{\partial L}{\partial \mathbf{v}}$ . On the other hand, the

equations  $\mathbf{P} = \frac{\partial L}{\partial \mathbf{v}}$  allow to express the velocity

$\mathbf{v}$  of an arbitrary particle through the generalized momentum  $\mathbf{P}$ . Then, we can substitute these velocities in (49) and determine  $H$  only through  $\mathbf{P}$ .

In order to find  $\frac{\partial L}{\partial \mathbf{v}}$  in (50), in each

particular Lagrangian  $L$ , we should express  $J^\mu$  and  $j^\mu$  in terms of the velocity  $\mathbf{v}$  and interval  $ds$ :

$$\begin{aligned} J^\mu &= c \rho_0 \frac{dx^\mu}{ds} = c \rho_0 \frac{dt}{ds} \frac{dx^\mu}{dt}, \\ j^\mu &= c \rho_{0q} \frac{dt}{ds} \frac{dx^\mu}{dt}, \end{aligned} \quad (51)$$

while  $dx^\mu = (cdt, d\mathbf{r})$ , and we introduce the

notation  $\frac{dx^\mu}{dt} = (c, \frac{d\mathbf{r}}{dt}) = (c, \mathbf{v}) = (c, v^i)$ , where

the four-dimensional quantity  $\frac{dx^\mu}{dt}$  is not a real four-vector. With regard to the definition of the

$$\begin{aligned} L &= \sum_{n=1}^{N_p} \int \left( \begin{aligned} &-\rho_0 \left( \mathcal{G} - \mathbf{v} \cdot \mathbf{U} \right) - \rho_0 (\psi - \mathbf{v} \cdot \mathbf{D}) \\ &-\rho_{0q} (\phi - \mathbf{v} \cdot \mathbf{A}) - \rho_0 (\varphi - \mathbf{v} \cdot \mathbf{\Pi}) \end{aligned} \right) \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0 \\ &+ \int \left( \begin{aligned} &c k R - 2c k \Lambda + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \\ &-\frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \end{aligned} \right) \sqrt{-g} dx^1 dx^2 dx^3, \end{aligned} \quad (54)$$

four-potential of the acceleration field

$u_\mu = \left( \frac{\mathcal{G}}{c}, -\mathbf{U} \right)$ , for each particle we obtain:

$$u_\mu J^\mu = c \rho_0 \frac{dt}{ds} \left( \mathcal{G} - \mathbf{v} \cdot \mathbf{U} \right). \quad (52)$$

In (48), the unit of volume of the system in any particular integral can be expressed in terms of the unit of volume in the reference frame  $K_p$  associated with the particle in the following way:

$$\sqrt{-g} dx^1 dx^2 dx^3 = \frac{ds}{cdt} \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0. \quad (53)$$

From this formula in the weak-field limit in Minkowski space, when  $ds = cdt/\gamma$ , it follows that the volume of a moving particle is decreased in comparison with the volume of a particle at rest. Given that  $ds = c d\tau$ , where  $\tau$  is the proper time in the reference frame  $K_p$  of the particle, the equality of four-volumes in different reference frames follows from (53):

$$\begin{aligned} \int \sqrt{-g} c dt dx^1 dx^2 dx^3 &= \\ \int \left( \sqrt{-g} c d\tau dx^1 dx^2 dx^3 \right)_0. \end{aligned}$$

This equation reflects the fact that the four-volume is a four-invariant.

Under the above conditions, (40), (51), (52) and (53) can be written for the Lagrangian (48) as follows:

as well as after partial volume integration:

$$L = - \sum_{n=1}^{N_p} \left( \begin{aligned} & m \left( \mathcal{G} - \mathbf{v} \cdot \mathbf{U} \right) + m \left( \psi - \mathbf{v} \cdot \mathbf{D} \right) \\ & + q \left( \varphi - \mathbf{v} \cdot \mathbf{A} \right) + m \left( \wp - \mathbf{v} \cdot \mathbf{\Pi} \right) \end{aligned} \right) + \int \left( \begin{aligned} & c k R - 2 c k \Lambda + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \\ & - \frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \end{aligned} \right) \sqrt{-g} dx^1 dx^2 dx^3, \quad (55)$$

where  $m = \int \rho_0 \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0$  is the mass

of an arbitrary particle,

$q = \int \rho_{0q} \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0$  is the particle's

charge. In (55), the scalar and vector field potentials are averaged over the particle's volume, which means that they are the effective potentials at the location of the particle.

In operations with three-vectors, it is convenient to write vectors in the form of components or projections on the spatial axes of the coordinate system using, for example, instead of the velocity  $\mathbf{v}$ , the quantity  $v^i$ , where  $i=1,2,3$ . Then,  $v^1 = v_x$ ,  $v^2 = v_y$ ,  $v^3 = v_z$ , and the velocity derivative can be represented as:  $\frac{\partial}{\partial \mathbf{v}} = \frac{\partial}{\partial v^i}$ . For the gravitational vector potential in particular, we obtain:  $\mathbf{D} = (D_x, D_y, D_z) = (D_1, D_2, D_3)$ .

With this in mind, from (55) and (50), we find:

$$H = \sum_{n=1}^{N_p} \left( \begin{aligned} & m \mathcal{G} + m \psi + q \varphi + m \wp \\ & - \int \left( c k R - 2 c k \Lambda + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \right. \\ & \left. - \frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} dx^1 dx^2 dx^3. \end{aligned} \right) \quad (58)$$

In (58), the Hamiltonian contains the scalar curvature  $R$  and the cosmological constant  $\Lambda$ . As it will be shown in Section 6 about the energy, this Hamiltonian represents the relativistic energy of the system. To make the

$$\begin{aligned} \mathbf{P} &= P_i = \sum_{n=1}^{N_p} \frac{\partial L}{\partial v^i} \\ &= \sum_{n=1}^{N_p} \left( m U_i + m D_i + q A_i + m \Pi_i \right), \\ \mathbf{P} &= m U_i + m D_i + q A_i + m \Pi_i. \end{aligned} \quad (56)$$

Based on this, we find for the sums of the scalar products of three-vectors by summing over the index  $i$ :

$$\sum_{n=1}^{N_p} \left( \begin{aligned} & \mathbf{P} \cdot \mathbf{v} = \\ & \sum_{n=1}^{N_p} \left( m U_i v^i + m D_i v^i + q A_i v^i + m \Pi_i v^i \right) \end{aligned} \right) \quad (57)$$

From (49), taking (55) and (57) into account, we have:

picture complete, we could also express the quantity  $\mathcal{G} = c g_{0\mu} u^\mu$  in (58) through the generalized momentum  $P_i$ . We have described this procedure in [4].

For continuously distributed matter, the masses and charges of the particles in (58) can be expressed through the corresponding integrals:

$$m = \int_0^n \rho_0 \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0,$$

$$q = \int_0^n \rho_{0q} \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0. \text{ Also, taking (53)}$$

into account, in which we can substitute the

$$H = \frac{1}{c} \int \left( \rho_0 \vartheta + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3 - \int \left( c k R - 2 c k \Lambda + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \left( -\frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} dx^1 dx^2 dx^3 \right). \quad (59)$$

### Hamilton's equations

Assuming that the Hamiltonian depends on the generalized three-momenta of particles  $\mathbf{P}^n$ :  $H = H(t, \mathbf{r}, \mathbf{P}, D_\mu, A_\mu, \pi_\mu)$  and the Lagrangian depends on three-velocity of particles  $\mathbf{v}^n$ :  $L = L(t, \mathbf{r}, \mathbf{v}, D_\mu, A_\mu, \pi_\mu)$ , where  $\mathbf{r}^n = (x, y, z)$  is a three-dimensional radius-vector of the particle with the number  $n$ , we will take differentials of  $L$  and  $H$ , as well as the differentials of both sides of equation (49):

$$DL = \frac{\partial L}{\partial t} Dt + \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{r}^n} D\mathbf{r}^n + \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{v}^n} D\mathbf{v}^n + \frac{\partial L}{\partial D_\mu} DD_\mu + \frac{\partial L}{\partial A_\mu} DA_\mu + \frac{\partial L}{\partial \pi_\mu} D\pi_\mu. \quad (60)$$

$$DH = \frac{\partial H}{\partial t} Dt + \sum_{n=1}^{N_p} \frac{\partial H}{\partial \mathbf{r}^n} D\mathbf{r}^n + \sum_{n=1}^{N_p} \frac{\partial H}{\partial \mathbf{P}^n} D\mathbf{P}^n + \frac{\partial H}{\partial D_\mu} DD_\mu + \frac{\partial H}{\partial A_\mu} DA_\mu + \frac{\partial H}{\partial \pi_\mu} D\pi_\mu. \quad (61)$$

$$DH = \sum_{n=1}^{N_p} \left( D\mathbf{P}^n \cdot \mathbf{v}^n \right) + \sum_{n=1}^{N_p} \left( \mathbf{P}^n \cdot D\mathbf{v}^n \right) - DL. \quad (62)$$

Substituting (60) and (61) into (62), we find:

expression  $\frac{cdt}{ds} = \frac{u^0}{c}$ , where  $u^0$  denotes the time component of the four-velocity of an arbitrary particle, from (58) we find:

$$\begin{aligned} \frac{\partial H}{\partial t} &= -\frac{\partial L}{\partial t}, \quad \frac{\partial H}{\partial \mathbf{r}^n} = -\frac{\partial L}{\partial \mathbf{r}^n}, \quad \frac{\partial H}{\partial D_\mu} = -\frac{\partial L}{\partial D_\mu}, \\ \frac{\partial H}{\partial A_\mu} &= -\frac{\partial L}{\partial A_\mu}, \quad \frac{\partial H}{\partial \pi_\mu} = -\frac{\partial L}{\partial \pi_\mu}, \quad \frac{\partial H}{\partial \mathbf{P}^n} = \mathbf{v}^n, \\ \frac{\partial L}{\partial \mathbf{v}^n} &= \mathbf{P}^n. \end{aligned} \quad (63)$$

The last equation in (63) leads to (50) and gives the expression (56) for the generalized momentum  $\mathbf{P}^n$  of an arbitrary particle of the system in an explicit form.

We will now apply the principle of least action to the Lagrangian in the form

$L = L(t, \mathbf{r}, \mathbf{v}, D_\mu, A_\mu, \pi_\mu)$ , equating the action variation to zero, when the particle moves from the time point  $t_1$  to the time point  $t_2$ .

$$\delta S = \delta \int_{t_1}^{t_2} L(t, \mathbf{r}, \mathbf{v}, D_\mu, A_\mu, \pi_\mu) dt + \int_{t_1}^{t_2} \left( \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{r}^n} \delta \mathbf{r}^n + \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{v}^n} \delta \mathbf{v}^n + \frac{\partial L}{\partial D_\mu} \delta D_\mu + \frac{\partial L}{\partial A_\mu} \delta A_\mu + \frac{\partial L}{\partial \pi_\mu} \delta \pi_\mu \right) dt = 0. \quad (64)$$

In (64), it was assumed that the time variation is equal to zero:  $\delta t = 0$ . Partial derivatives with variations  $\delta D_\mu$ ,  $\delta A_\mu$  and  $\delta \pi_\mu$  lead to field equations (16), (18) and (26). If we take into account the definition of velocity in the second

term in the integral (64):  $\mathbf{v} = \frac{d\mathbf{r}}{dt}$ , then the integral for this term is taken by parts. Then, for the first and second terms in the integral (64), we have the following:

$$\left. \begin{aligned} \sum_{n=1}^{N_p} \int_{t_1}^{t_2} \left( \frac{\partial L}{\partial \mathbf{r}} \delta \mathbf{r} + \frac{\partial L}{\partial \mathbf{v}} d \delta \mathbf{r} \right) = \\ \sum_{n=1}^{N_p} \frac{\partial L}{\partial \mathbf{v}} \delta \mathbf{r} \Big|_{t_1}^{t_2} + \sum_{n=1}^{N_p} \int_{t_1}^{t_2} \left( \frac{\partial L}{\partial \mathbf{r}} - \frac{d}{dt} \frac{\partial L}{\partial \mathbf{v}} \right) \delta \mathbf{r} dt. \end{aligned} \right\} \quad (65)$$

When varying the action, the variations  $\delta \mathbf{r}$  are equal to zero only at the beginning and at the end of the motion; that is at  $t_1$  and  $t_2$ . Therefore, for vanishing of the variation  $\delta S$ , it is necessary that the quantity in brackets inside the integral (65) would be equal to zero. This leads to the well-known Lagrange equations of motion:

$$\frac{\partial L}{\partial \mathbf{r}} = \frac{d}{dt} \frac{\partial L}{\partial \mathbf{v}}. \quad (66)$$

According to (63),  $\frac{\partial L}{\partial \mathbf{v}} = \mathbf{P}$ , as well as

$$\frac{\partial H}{\partial \mathbf{r}} = - \frac{\partial L}{\partial \mathbf{r}}. \text{ Let us substitute this in (66):}$$

$$\frac{\partial H}{\partial \mathbf{r}} = - \frac{d \mathbf{P}}{dt}. \quad (67)$$

Equation (67) together with equation  $\frac{\partial H}{\partial \mathbf{P}} = \mathbf{v}$  from (63) represent the standard

Hamiltonian equations describing the motion of an arbitrary particle of the system in the gravitational and electromagnetic fields and in the pressure field. According to (67), the rate of change of the generalized momentum of the particle by the coordinate time is equal to the generalized force, which is found as the gradient with respect to the particle's coordinates of the relativistic energy of the system taken with the

opposite sign. These equations are widely used; not only in the general theory of relativity, but also in other areas of theoretical physics. We have checked these equations in [4] in the framework of the covariant theory of gravitation by direct substitution of the Hamiltonian.

## The system's energy

We will consider a closed system which is in the state of some stationary motion. An example would be a charged ball rotating around its center of mass, which forms the system under consideration together with its gravitational and electromagnetic fields and the internal pressure. In such a system, the energy should be conserved as a consequence of lack of energy losses to the environment and taking into account the homogeneity of time; i.e., the equivalence of the time points for the system's state.

The system's Lagrangian, taking the fields' energy into account, has the form of (55). Due to the stationary motion, we can assume that within the system's volume the metric tensor  $g_{\mu\nu}$ , the scalar curvature  $R$ , the four-potentials of the field  $D_\mu$ ,  $A_\mu$  and of the pressure  $\pi_\mu$  do not depend on time. But, since any point particle moves with the ball, then its location and velocity are changed, being defined by the radius vector  $\mathbf{r}$  and velocity  $\mathbf{v}$ , respectively. We may assume that the Lagrangian of the system does not depend explicitly on time and is a function of

the form:  $L = L(\mathbf{r}, \mathbf{v})$ . Now, we will take the time derivative of the Lagrangian, as it is done for example in [13], not only for one but for a set of particles, and will apply (66):

$$\begin{aligned} \frac{dL}{dt} &= \frac{1}{dt} \sum_{n=1}^{N_p} \left( \frac{\partial L}{\partial \mathbf{r}} d \mathbf{r} + \frac{\partial L}{\partial \mathbf{v}} d \mathbf{v} \right) \\ &= \sum_{n=1}^{N_p} \left( \frac{\partial L}{\partial \mathbf{r}} \mathbf{v} + \frac{\partial L}{\partial \mathbf{v}} \frac{d \mathbf{v}}{dt} \right) \\ &= \sum_{n=1}^{N_p} \left( \mathbf{v} \frac{d}{dt} \frac{\partial L}{\partial \mathbf{v}} + \frac{\partial L}{\partial \mathbf{v}} \frac{d \mathbf{v}}{dt} \right) \\ &= \frac{d}{dt} \sum_{n=1}^{N_p} \left( \mathbf{v} \frac{\partial L}{\partial \mathbf{v}} \right). \end{aligned}$$



$$\frac{d}{dt} \left( \sum_{n=1}^{N_p} \left( \mathbf{v} \frac{\partial L}{\partial \mathbf{v}} \right) - L \right) = 0.$$

The quantity in the brackets is not time-dependent and is constant. This gives the definition of relativistic energy as a conserved quantity for a closed system at stationary motion:

$$E = \sum_{n=1}^{N_p} \left( \mathbf{v} \frac{\partial L}{\partial \mathbf{v}} \right) - L. \quad (68)$$

With regard to (63) and (49), we find the following:

$$E = \frac{1}{c} \int \left( \rho_0 \vartheta + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3 - \int \left( c k R - 2c k \Lambda + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \left( -\frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} dx^1 dx^2 dx^3 \right). \quad (70)$$

Using expression (70), we can find the invariant energy  $E_0$  of the system, for which we should use the frame of reference of the center of mass and calculate the integral. In addition, at a known velocity  $\mathbf{v}$  of the center of mass of the system in an arbitrary reference frame  $K$ , we can calculate the momentum of the system in  $K$ . This can be clarified as follows. We will define the invariant mass of the system taking the mass-energy of the fields into account using the relation:  $m_0 = \frac{E_0}{c^2}$ , where  $c$  is the speed of

light as a measure of the velocity of propagation of electromagnetic and gravitational interactions. If the four-displacement in  $K$  has the form:  $d\hat{x}^\mu = (cdt, dx, dy, dz) = (cdt, d\mathbf{r}) = dt(c, \mathbf{v})$ , then, for the four-velocity of the system in  $K$ , we can write:  $\hat{u}^\mu = \frac{cd\hat{x}^\mu}{ds} = \frac{cdt}{ds} \frac{d\hat{x}^\mu}{dt} = \frac{cdt}{ds} (c, \mathbf{v})$ . The four-vector  $p^\mu = m_0 \hat{u}^\mu = \frac{cdt}{ds} (m_0 c, m_0 \mathbf{v}) = \left( \frac{E}{c}, \mathbf{p} \right)$  defines the four-momentum, which contains the relativistic

$$E = \sum_{n=1}^{N_p} \left( \mathbf{P} \cdot \mathbf{v} \right) - L = H. \quad (69)$$

It turns out that the relativistic energy can be expressed in a covariant form, since according to (69) the formula for the energy coincides with the formula for the Hamiltonian in (49).

To calculate the relativistic energy of the system with the matter, which is continuously distributed over the volume, it is convenient to pass from the mass and charge of the particle to the corresponding densities inside the particle. According to (59), we obtain:

energy  $E = \frac{cdt}{ds} m_0 c^2 = \frac{cdt}{ds} E_0$  and relativistic momentum  $\mathbf{p} = \frac{cdt}{ds} m_0 \mathbf{v} = \frac{cdt}{ds} \frac{E_0}{c^2} \mathbf{v}$ . This gives the formula for determining the momentum through the energy:  $\mathbf{p} = \frac{E}{c^2} \mathbf{v}$ , and, correspondingly, for the four-momentum:  $p^\mu = \left( \frac{E}{c}, \mathbf{p} \right) = \left( \frac{E}{c}, \frac{E}{c^2} \mathbf{v} \right)$ .

In the reference frame  $K'$ , in which the system is at rest  $\mathbf{v} = 0$ ,  $\mathbf{p} = 0$ ,  $dt = d\tau$ , and then  $E = E_0$ , and also  $(p^\mu)_{\mathbf{v}=0} = \left( \frac{E_0}{c}, 0, 0, 0 \right)$ ; that is in the four-momentum in the reference frame  $K'$ , only the time component is non-zero.

If we multiply the four-momentum by the speed of light, we will obtain the four-vector of the form  $H^\mu = c p^\mu = (E, c\mathbf{p}) = \left( E, \frac{E}{c} \mathbf{v} \right)$ , the time component of which is the relativistic energy, equal in value to the Hamiltonian. Thus, we find the four-vector, which in [4] was called the Hamiltonian four-vector.

## The Cosmological Constant Gauge and the Resulting Consequences

We will make transformations and substitute (43) and (39) in (70):

$$E = \frac{1}{c} \int \left( \rho_0 \mathcal{G} + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3 - \int \left( \frac{2ck\Lambda + 2\chi + \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu}}{-\frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu}} \right) \sqrt{-g} dx^1 dx^2 dx^3. \quad (71)$$

If we choose the condition for the cosmological constant in the form:

$$2ck\Lambda + 2\chi = 0, \quad (72)$$

then the relativistic energy (71) is uniquely defined, since the dependence on the constants  $\Lambda$  and  $\chi$  disappears:

$$E = \frac{1}{c} \int \left( \rho_0 \mathcal{G} + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3 - \int \left( \frac{\frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu}}{-\frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu}} \right) \sqrt{-g} dx^1 dx^2 dx^3. \quad (73)$$

We will remind that the quantities  $\Lambda$  and  $\chi$  can have their own values for each particle of matter. But on condition of (72), the expression for the relativistic energy (73) becomes universal for any particle in an arbitrary system of particles and their fields.

From (72) and (39), the following equation is obtained:

$$\left. \begin{aligned} 2ck\Lambda &= -2D_\mu J^\mu - 2A_\mu j^\mu \\ &- 2u_\mu J^\mu - 2\pi_\mu J^\mu. \end{aligned} \right\} \quad (74)$$

In order to estimate the value of the cosmological constant  $\Lambda$ , it is convenient to divide all of the system's matter into small pieces, scatter them apart to infinity and leave them motionless. Then, the vector potentials of the fields and pressure become equal to zero and the relation remains:

$$ck\Lambda = (-\rho_0 \psi_0 - \rho_{0q} \varphi_0 - \rho_0 c^2 - p_0)_\infty.$$

It follows that  $\Lambda$ , just like  $\chi$  in (42), is associated with the rest energy, with the pressure energy and with the proper energy of the fields of the system under consideration.

If in some volume there are no particles and the mass density  $\rho_0$  and the charge density  $\rho_{0q}$

are zero, then in this volume there must remain the relativistic energy of the external fields:

$$E_{rf} = - \int \left( \frac{\frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu}}{-\frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu}} \right) \sqrt{-g} dx^1 dx^2 dx^3. \quad (75)$$

Based on (73), we can express the energy of a small body at rest. For simplicity, we will assume that the body does not rotate as a whole and there is no motion of the matter and charges inside it (an ideal solid body without the intrinsic magnetic field and the torsion field). Under such conditions, the coordinate time of the system becomes approximately equal to the proper time of the body:  $dt \approx d\tau$ . Since the interval  $ds = c d\tau$ , then we obtain:  $ds \approx c dt$ . Since there is no spatial motion in any part of the body, we can write:

$$\begin{aligned} \mathcal{G} &= c g_{0\mu} u^\mu = g_{0\mu} \frac{c^2 dx^\mu}{ds} \\ &= g_{0\mu} \frac{c^2}{ds} (c dt, 0, 0, 0) = g_{00} \frac{c^3 dt}{ds} = c^2 g_{00}. \\ u^0 &= \frac{c dx^0}{ds} \approx c. \end{aligned}$$

With this in mind, we obtain from (73):

$$E_0 = \int \left( g_{00} \rho_0 c^2 + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) \sqrt{-g} dx^1 dx^2 dx^3 - \int \left( \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} dx^1 dx^2 dx^3. \quad (76)$$

In the weak field limit in (76), we can use  $g_{00} \approx 1$ ,  $\sqrt{-g} \approx 1$ . The tensor product  $u_{\mu\nu} u^{\mu\nu}$  in the absence of matter motion inside the ideal solid body vanishes. Using (F5) and (F6), we can write:

$$\begin{aligned} \mathbf{C} &= -\nabla \left( \frac{p_0}{\rho_0} \right), \quad \mathbf{I} = 0, \\ \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} &= -\frac{1}{8\pi\sigma} (C^2 - c^2 I^2) \\ &= -\frac{1}{8\pi\sigma} \left[ \nabla \left( \frac{p_0}{\rho_0} \right) \right]^2. \end{aligned}$$

Besides, in [4] it was found that in the weak field for a motionless body in the form of a ball with uniform density of mass and charge, the following relations hold for the body's proper fields:

$$\begin{aligned} -\frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} &= \frac{1}{8\pi G} \Gamma^2, \\ \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} &= -\frac{\varepsilon_0}{2} E^2, \\ \int \left( \frac{1}{2} \rho_0 \psi \right) dx^1 dx^2 dx^3 &= \\ \int \left( -\frac{1}{8\pi G} \Gamma^2 \right) dx^1 dx^2 dx^3, \\ \int \left( \frac{1}{2} \rho_{0q} \varphi \right) dx^1 dx^2 dx^3 &= \\ \int \left( \frac{\varepsilon_0}{2} E^2 \right) dx^1 dx^2 dx^3. \end{aligned} \quad (77)$$

According to (77), the potential energy of the ball's matter in the proper gravitational field which is associated with the scalar potential  $\psi$  is twice the potential energy associated with the

field strength  $\Gamma$ . The same is true for the electromagnetic field with the potential  $\varphi$  and the strength  $\mathbf{E}$  both in the case of uniform arrangement of charges in the ball's volume and in case of their location on the surface only. Substituting (77) into (76) in the framework of the special theory of relativity gives the invariant energy of the system in the form of a fixed solid spherical body with uniform density of mass and charge, taking into account the energy of their proper potential fields:

$$(E_0)_{SR} = \int \left( \rho_0 c^2 + \frac{1}{2} \rho_0 \psi + \frac{1}{2} \rho_{0q} \varphi + p_0 - \frac{1}{8\pi\sigma} \left[ \nabla \left( \frac{p_0}{\rho_0} \right) \right]^2 \right) dx^1 dx^2 dx^3. \quad (78)$$

This calculation is apparently incomplete, since in reality inside any body there are particles, which cannot be as motionless as the body itself is. Therefore, in (78), in addition to the pressure and its gradient within the body, it is necessary to add the kinetic energy of motion of all the particles which constitute the body.

### The metric

Substituting (74) and (39) into (43), we find the expression for the scalar curvature  $R$ :

$$\begin{aligned} ckR &= -2D_\mu J^\mu - 2A_\mu j^\mu \\ &\quad - 2u_\mu J^\mu - 2\pi_\mu J^\mu \\ &= -2\chi, \end{aligned} \quad (79)$$

while  $ck = -\frac{c^4}{16\pi G\beta}$ , where  $\beta$  is a constant of the order of unity.

As it can be seen, the scalar curvature is zero in the whole space outside the body. The equation  $R=0$  does not mean, however, that the spacetime is flat as in the special theory of relativity, since the curvature of spacetime is determined by the components of the Riemann curvature tensor.

We will now substitute (74) into the equation for the metric (38):

$$\left. \begin{aligned} -2ck R^{\alpha\beta} + ck R g^{\alpha\beta} = \\ -D_\mu J^\mu g^{\alpha\beta} + U^{\alpha\beta} - A_\mu j^\mu g^{\alpha\beta} + W^{\alpha\beta} \\ -u_\mu J^\mu g^{\alpha\beta} + B^{\alpha\beta} - \pi_\mu J^\mu g^{\alpha\beta} + P^{\alpha\beta}, \end{aligned} \right\} \quad (80)$$

that also can be written using (39) as follows:

$$\left. \begin{aligned} -2ck R^{\alpha\beta} + ck R g^{\alpha\beta} = \\ U^{\alpha\beta} + W^{\alpha\beta} + B^{\alpha\beta} + P^{\alpha\beta} - \chi g^{\alpha\beta}. \end{aligned} \right\} \quad (81)$$

If we take the covariant derivative of (81), the left side of the equation vanishes due to the property of the Einstein tensor located there. The right side, with regard to the equation of motion (35) and provided that the metric tensor  $g^{\alpha\beta}$  in covariant differentiation behaves as a constant and  $\chi$  is a constant, vanishes too.

In (80), we can use (79) to replace the scalar curvature:

$$\left. \begin{aligned} -2ck R^{\alpha\beta} = D_\mu J^\mu g^{\alpha\beta} + U^{\alpha\beta} \\ + A_\mu j^\mu g^{\alpha\beta} + W^{\alpha\beta} \\ + u_\mu J^\mu g^{\alpha\beta} + B^{\alpha\beta} \\ + \pi_\mu J^\mu g^{\alpha\beta} + P^{\alpha\beta}. \end{aligned} \right\} \quad (82)$$

If we sum up (80) and (82) and divide the result by 2, we will obtain the following equation for the metric:

$$\left. \begin{aligned} R^{\alpha\beta} - \frac{1}{4} R g^{\alpha\beta} = \\ -\frac{1}{2ck} (B^{\alpha\beta} + U^{\alpha\beta} + W^{\alpha\beta} + P^{\alpha\beta}), \end{aligned} \right\} \quad (83)$$

while with regard to (79)  $\nabla_\beta R = 0$ , according to (35)  $\nabla_\beta (B^{\alpha\beta} + U^{\alpha\beta} + W^{\alpha\beta} + P^{\alpha\beta}) = 0$  and

$\nabla_\beta \left( R^{\alpha\beta} - \frac{1}{2} R g^{\alpha\beta} \right) = 0$  as the property of the Einstein tensor.

In empty space according to (83), the curvature tensor  $R^{\alpha\beta}$  depends only on the stress-energy tensors of the gravitational and electromagnetic fields  $U^{\alpha\beta}$  and  $W^{\alpha\beta}$ , so these fields change the curvature of spacetime outside the bodies. We will note that in equation (83), the cosmological constant  $\Lambda$  and the tensor product of the type  $D_\mu J^\mu g^{\alpha\beta}$  are missing. This fact makes determination of the metric tensor components much easier.

If we compare (83) with the Einstein equation (46), then two major differences will be found out — at the right side of (83) stress-energy tensors  $U^{\alpha\beta}$ ,  $B^{\alpha\beta}$  and  $P^{\alpha\beta}$  are present, and in addition the coefficient in front of the scalar curvature  $R$  is two times less than in (46).

## The Energy Components

In Newtonian mechanics, the relations for the Lagrangian and the total energy are known:  $L = E_k - U$ ,  $E_t = E_k + U$ , where  $E_k$  denotes the kinetic energy, which depends only on the velocity, and  $U$  denotes the potential energy of the system, depending both on the coordinates and the velocity. In relativistic physics, instead of individual scalar functions and three-dimensional vectors, four-vectors and four-tensors are used, in which the scalar functions and three-dimensional vectors are combined into one whole. In addition, instead of the negative total energy  $E_t$ , the positive relativistic energy

$E$  is usually used. While we have already determined the energy  $E$  in (73), then for the Lagrangian (54) we should additionally replace the scalar curvature  $R$  with the help of (79) and the cosmological constant  $\Lambda$  with the help of (74). Taking the relation:  $\left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0$

$= \frac{u^0}{c} \sqrt{-g} dx^1 dx^2 dx^3$  into account this will give the following:

$$L = \frac{1}{c} \sum_{n=1}^{N_p} \int^n \left\{ \begin{aligned} & -\rho_0 \left( \mathcal{G} - \mathbf{v} \cdot \mathbf{U} \right) - \rho_0 (\psi - \mathbf{v} \cdot \mathbf{D}) \\ & -\rho_{0q} (\varphi - \mathbf{v} \cdot \mathbf{A}) - \rho_0 (\wp - \mathbf{v} \cdot \mathbf{\Pi}) \end{aligned} \right\} u^0 \sqrt{-g} dx^1 dx^2 dx^3 \\ + \int \left\{ \begin{aligned} & \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} \\ & -\frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \end{aligned} \right\} \sqrt{-g} dx^1 dx^2 dx^3, \quad (84)$$

Calculating the energy  $E_V$ , which is associated with the four-dimensional motion, as a half-sum of the relativistic energy (73) and the Lagrangian (84), we find:

$$E_V = \frac{1}{2} (E + L) \\ = \frac{1}{2c} \sum_{n=1}^{N_p} \int^n \left( \rho_0 \mathbf{v} \cdot \mathbf{U} + \rho_0 \mathbf{v} \cdot \mathbf{D} + \rho_{0q} \mathbf{v} \cdot \mathbf{A} + \rho_0 \mathbf{v} \cdot \mathbf{\Pi} \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3. \quad (85)$$

where  $\mathbf{v}$  is three-velocity vector of the particle with the number  $n$ .

If in (57) we replace the masses  $m$  and charges  $q$  by the corresponding integrals in the form:

$$m = \int^n \rho_0 \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0,$$

$$q = \int^n \rho_{0q} \left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0, \text{ and transform}$$

the volume units in the form

$$\left( \sqrt{-g} dx^1 dx^2 dx^3 \right)_0 = \frac{u^0}{c} \sqrt{-g} dx^1 dx^2 dx^3, \text{ then}$$

we obtain the relation  $E_V = \frac{1}{2} \sum_{n=1}^{N_p} \left( \mathbf{P} \cdot \mathbf{v} \right)$ . As we

can see, the kinetic energy  $E_V$  of the system in the reference frame of the center of mass vanishes only when the velocities  $\mathbf{v}$  of all the system's particles at the same time vanish.

We will determine the potential energy  $U_P$  as a half-difference of the relativistic energy (73) and the Lagrangian (84):

$$U_P = \frac{1}{2} (E - L) = \frac{1}{c} \int \left( \rho_0 \mathcal{G} + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3 \\ - \frac{1}{2c} \sum_{n=1}^{N_p} \int^n \left( \rho_0 \mathbf{v} \cdot \mathbf{U} + \rho_0 \mathbf{v} \cdot \mathbf{D} + \rho_{0q} \mathbf{v} \cdot \mathbf{A} + \rho_0 \mathbf{v} \cdot \mathbf{\Pi} \right) u^0 \sqrt{-g} dx^1 dx^2 dx^3 \\ - \int \left( \frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} - \frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} - \frac{c^2}{16\pi\eta} u_{\mu\nu} u^{\mu\nu} - \frac{c^2}{16\pi\sigma} f_{\mu\nu} f^{\mu\nu} \right) \sqrt{-g} dx^1 dx^2 dx^3. \quad (86)$$

For a solid body in the limit of the special theory of relativity, when  $\sqrt{-g} = 1$ ,  $u^0 = c\gamma$ ,

$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$ , the expression for the kinetic energy (85) of the system is as follows:

$$(E_V)_{SR} = \frac{1}{2} \sum_{n=1}^{N_p} \int \left( \rho_0 \mathbf{v} \cdot \mathbf{U} + \rho_0 \mathbf{v} \cdot \mathbf{D} + \rho_{0q} \mathbf{v} \cdot \mathbf{A} + \rho_0 \mathbf{v} \cdot \mathbf{\Pi} \right) \gamma dx^1 dx^2 dx^3. \quad (87)$$

The main part of the kinetic energy is proportional to the square of the velocity, and vector potentials of all fields, including the velocity field and pressure field, make contribution to this part of the energy.

For the potential energy (86) of a solid body in the limit of special theory of relativity, the tensor product  $u_{\mu\nu} u^{\mu\nu}$  according to (E8) tends to zero. We will also take into account the values of other tensor products:

$$\frac{c^2}{16\pi G} \Phi_{\mu\nu} \Phi^{\mu\nu} = -\frac{1}{8\pi G} (\Gamma^2 - c^2 \Omega^2),$$

$$\frac{c^2 \varepsilon_0}{4} F_{\mu\nu} F^{\mu\nu} = -\frac{\varepsilon_0}{2} (E^2 - c^2 B^2),$$

$$\frac{c^2}{16\pi \sigma} f_{\mu\nu} f^{\mu\nu} = -\frac{1}{8\pi \sigma} (C^2 - c^2 I^2).$$

It gives the following:

$$\begin{aligned} (U_P)_{SR} = & \int \left( \rho_0 \wp + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) \gamma dx^1 dx^2 dx^3 \\ & - \frac{1}{2} \sum_{n=1}^{N_p} \int \left( \rho_0 \mathbf{v} \cdot \mathbf{U} + \rho_0 \mathbf{v} \cdot \mathbf{D} + \rho_{0q} \mathbf{v} \cdot \mathbf{A} + \rho_0 \mathbf{v} \cdot \mathbf{\Pi} \right) \gamma dx^1 dx^2 dx^3 \\ & + \int \left( \frac{1}{8\pi G} (\Gamma^2 - c^2 \Omega^2) - \frac{\varepsilon_0}{2} (E^2 - c^2 B^2) - \frac{1}{8\pi \sigma} (C^2 - c^2 I^2) \right) dx^1 dx^2 dx^3. \end{aligned}$$

The potential energy depends also on the velocity. If  $\mathbf{v} = 0$  for all material particles of the

system, then the potential energy of the system remains, taking the field energy into account:

$$\begin{aligned} (U_P)_{SR} = & \int \left( \rho_0 c^2 + \rho_0 \psi + \rho_{0q} \varphi + \rho_0 \wp \right) dx^1 dx^2 dx^3 \\ & + \int \left( \frac{1}{8\pi G} (\Gamma^2 - c^2 \Omega^2) - \frac{\varepsilon_0}{2} (E^2 - c^2 B^2) - \frac{1}{8\pi \sigma} (C^2 - c^2 I^2) \right) dx^1 dx^2 dx^3. \end{aligned} \quad (88)$$

In the absence of external fields and internal motions in the fixed system, in (88) the fields  $\Omega$ ,  $B$ ,  $I$  become equal to zero. As a result, with regard to (77), the potential energy becomes equal to the relativistic energy (78) for a fixed ideal solid body.

## Conclusions

We have presented the Lagrangian of the system as consisting of one term for the curvature and four pairs of terms of identical form for each of the four fields: gravitational and electromagnetic fields, acceleration field and pressure field. As a result, for each field we have obtained equations coinciding in form with each other. The spacetime is also represented by its proper tensor metric field  $g_{\alpha\beta}$ . The mass four-current  $J^\mu$  interacts with the specified fields, gaining energy in them. Moreover, the

electromagnetic field changes the energy of electromagnetic four-current  $j^\mu$ . However, the fields also have their proper energy and momentum, which are part of the tensors  $U^{\alpha\beta}$  (5),  $W^{\alpha\beta}$  (8),  $B^{\alpha\beta}$  (11) and  $P^{\alpha\beta}$  (14), respectively.

The similarity of field equations implies the necessity of gauge; not only of the four-potentials of the gravitational and electromagnetic fields, but also of the four-potentials of the acceleration field and pressure field, as well as of the mass four-current  $J^\mu$  and electromagnetic four-current  $j^\mu$ . From the standpoint of physics, the meaning of such gauges is that the source of divergence of the three-velocity vector of small volume may be the time changes in the particle's energy in any fields which are present in the given volume.

This may be the particle's energy in the velocity field, the energy in the pressure field or the energy in the gravitational or electromagnetic fields.

In contrast to the standard approach, we do not use any of the variety of known forms of stress-energy tensors of matter. Instead, the energy and momentum of the matter are described based on the acceleration field, the acceleration field tensor and the stress-energy tensor of the acceleration field. The contribution of pressure to the system's energy and momentum, respectively, is described through the pressure field with the help of the pressure field tensor and the stress-energy tensor of the pressure field. In this case, the acceleration field and the pressure field, as well as the electromagnetic and gravitational fields, are regarded as the four-dimensional vector fields with their own four-potentials.

Representation of the gravitational field as a vector field is performed within the covariant theory of gravitation [3-4], in contrast to the general theory of relativity, where gravitation is described indirectly through the spacetime geometry and is considered as a metric tensor field. We consider as an advantage of our approach the fact that the energy and momentum of the gravitational field at each point are uniquely determined with the help of the stress-energy tensor of the gravitational field; whereas, in the general theory of relativity, we have to restrict ourselves only to the corresponding pseudotensor, such as the Landau-Lifshitz stress-energy pseudotensor [13].

In order to uniquely identify the relativistic energy of a particle or a matter unit, we used a special gauge of the cosmological constant, giving this constant the meaning of the rest energy of the particle with "turned-off" external fields and influences. This led to the expression for the relativistic energy of the system (73) and to the equation for the metric (83), the right side of which is the sum of four stress-energy tensors of the fields.

In the absence of the cosmological constant in the Lagrangian (1), it would be impossible to perform the specified calibration, the physical system's energy would be uncertain and the presented theory would remain unfinished. In our approach, the cosmological constant does not reflect the energy density of the empty cosmic space, or the so-called dark energy, but rather the

energy density of the matter scattered in space.

Given that  $-2ck = \frac{c^4}{8\pi G\beta}$ , where  $\beta$  is a

constant of the order of unity, it follows from

(74) that  $\Lambda \approx \frac{16\pi G\rho_0\beta}{c^2}$ . Substituting here the

standard estimate of the cosmological constant  $\Lambda \approx 10^{-52} \text{ m}^{-2}$ , we find the corresponding mass density:  $\rho_0 \approx 3 \times 10^{-27} \text{ kg/m}$ . This density is sufficiently close to the density of cosmic matter, averaged over the entire space.

It can be noted that our expression for the relativistic energy and the equation for the metric differ substantially from those obtained in the general theory of relativity. For the energy it follows from the fact that instead of the stress-energy tensor of matter we use the stress-energy tensors of the acceleration field and the pressure field, while the gravitational field is directly included in the energy, and not indirectly through the metric.

Let us take the Einstein equation for the metric with the cosmological constant from [14]. In the general case, the right-hand side of this equation contains the stress-energy tensor of the electromagnetic field  $W^{\alpha\beta}$  and the stress-energy tensor of matter  $\phi^{\alpha\beta}$ :

$$\left. \begin{aligned} R^{\alpha\beta} - \frac{1}{2} R g^{\alpha\beta} + \Lambda g^{\alpha\beta} = \\ \frac{8\pi G}{c^4} (W^{\alpha\beta} + \phi^{\alpha\beta}). \end{aligned} \right\} \quad (89)$$

The simplest form of the stress-energy tensor of matter without taking the pressure into account is the expression in terms of the mass density and the four-velocity:  $\phi^{\alpha\beta} = \rho_0 u^\alpha u^\beta$ . Contraction (89) with the metric tensor  $g_{\alpha\beta}$  gives the following:

$$\Lambda = \frac{8\pi G\phi}{4c^4} + \frac{R}{4},$$

where  $\phi = g_{\alpha\beta} \phi^{\alpha\beta}$ .

After substituting  $\Lambda$  in (89), the equation for the metric is transformed as follows:

$$\left. \begin{aligned} R^{\alpha\beta} - \frac{1}{4} R g^{\alpha\beta} = \\ \frac{8\pi G}{c^4} \left( W^{\alpha\beta} + \phi^{\alpha\beta} - \frac{\phi}{4} g^{\alpha\beta} \right) \end{aligned} \right\} \quad (90)$$

Now, equation for the metric (90) in the general theory of relativity can be compared with our equation for the metric (83). The main difference is that in (83) all the tensors at the right side act the same way, and in contraction with the metric tensor they vanish. But, it is not the case in (90) – if the expression  $g_{\alpha\beta} W^{\alpha\beta} = 0$  is valid for the electromagnetic stress-energy tensor, then for the stress-energy tensor of matter

the relation  $g_{\alpha\beta} \phi^{\alpha\beta} = 0$  does not hold, so that

in (90) one more term  $-\frac{\phi}{4} g^{\alpha\beta}$  is needed. As a

result, in the general theory of relativity, not only the gravitational field is represented in a special way, through the metric tensor, which differs from the method of introducing the electromagnetic field into the equation for the metric, but also the stress-energy tensor of matter  $\phi^{\alpha\beta}$  is asymmetric with respect to the metric, in contrast to the stress-energy tensor of the electromagnetic field  $W^{\alpha\beta}$ .

### Appendix A. Variation of the sixth term in the action function

We need to find the variation for the sixth term in (1):

$$\delta S_6 = -\frac{1}{c} \int \delta \left( u_\mu J^\mu \sqrt{-g} \right) d\Sigma. \quad (A1)$$

We will need the expressions for variation of the metric tensor and the four-vector of the mass current, which can be found, for example, in [7], [9] and [14]:

$$\begin{aligned} \delta g^{\mu\nu} &= -g^{\mu\alpha} g^{\nu\beta} \delta g_{\alpha\beta}, \\ \delta \sqrt{-g} &= \frac{\sqrt{-g}}{2} g^{\alpha\beta} \delta g_{\alpha\beta}. \end{aligned} \quad (A2)$$

$$\left. \begin{aligned} \delta J^\beta &= \nabla_\sigma \left( J^\sigma \xi^\beta - J^\beta \xi^\sigma \right) \\ &= \frac{1}{\sqrt{-g}} \partial_\sigma \left[ \sqrt{-g} \left( J^\sigma \xi^\beta - J^\beta \xi^\sigma \right) \right] \end{aligned} \right\} \quad (A3)$$

In view of (A2) and (A3), we have the following:

$$\left. \begin{aligned} \delta \left( u_\mu J^\mu \sqrt{-g} \right) &= u_\mu \sqrt{-g} \delta J^\mu + u_\mu J^\mu \delta \sqrt{-g} \\ &\quad + J^\mu \sqrt{-g} \delta u_\mu \\ &= u_\mu \partial_\sigma \left[ \sqrt{-g} \left( J^\sigma \xi^\mu - J^\mu \xi^\sigma \right) \right] \\ &\quad + \frac{1}{2} u_\mu J^\mu g^{\alpha\beta} \sqrt{-g} \delta g_{\alpha\beta} \\ &\quad + J^\mu \sqrt{-g} \delta u_\mu. \end{aligned} \right\} \quad (A4)$$

We will transform the first term in (A4) with the help of functions' product differentiation by parts:

$$\begin{aligned} u_\mu \partial_\sigma \left[ \sqrt{-g} \left( J^\sigma \xi^\mu - J^\mu \xi^\sigma \right) \right] &= \\ \partial_\sigma \left[ u_\mu \sqrt{-g} \left( J^\sigma \xi^\mu - J^\mu \xi^\sigma \right) \right] &= \\ -\sqrt{-g} \left( J^\sigma \xi^\mu - J^\mu \xi^\sigma \right) \partial_\sigma u_\mu. \end{aligned}$$

In action variation, the term with the divergence can be neglected; the remaining term can be transformed as follows:

$$\begin{aligned} -\sqrt{-g} \left( J^\sigma \xi^\mu - J^\mu \xi^\sigma \right) \partial_\sigma u_\mu &= \\ -(\partial_\sigma u_\mu - \partial_\mu u_\sigma) J^\sigma \xi^\mu \sqrt{-g} &= \\ u_{\mu\sigma} J^\sigma \xi^\mu \sqrt{-g}. \end{aligned}$$

Substituting these results in (A4) and then in (A1), we find:

$$\delta S_6 = \int \left( \begin{aligned} &-\frac{1}{c} u_{\beta\sigma} J^\sigma \xi^\beta - \frac{1}{c} J^\beta \delta u_\beta \\ &-\frac{1}{2c} u_\mu J^\mu g^{\alpha\beta} \delta g_{\alpha\beta} \end{aligned} \right) \sqrt{-g} d\Sigma.$$



### Appendix B. Variation of the seventh term in the action function

Variation for the seventh term in (1) for the special case, when  $\eta$  is a constant, taking into account (A2) will be equal to:

$$\delta S_7 = -\frac{c}{16\pi\eta} \int \delta \left( u_{\mu\nu} u^{\mu\nu} \sqrt{-g} \right) d\Sigma, \quad (B1)$$

$$\left. \begin{aligned} & \delta \left( u_{\mu\nu} u^{\mu\nu} \sqrt{-g} \right) \\ &= \delta \left( u_{\mu\nu} u^{\mu\nu} \right) \sqrt{-g} + u_{\mu\nu} u^{\mu\nu} \delta \sqrt{-g} \\ &= u_{\mu\nu} \delta u^{\mu\nu} \sqrt{-g} + u^{\mu\nu} \delta u_{\mu\nu} \sqrt{-g} \\ &+ \frac{1}{2} u_{\mu\nu} u^{\mu\nu} g^{\alpha\beta} \sqrt{-g} \delta g_{\alpha\beta}. \end{aligned} \right\}$$

Since  $u^{\mu\nu} = g^{\mu\alpha} g^{\beta\nu} u_{\alpha\beta}$ , the tensor  $u_{\alpha\beta}$  is antisymmetric, then using the expression for  $\delta g^{\beta\nu}$  from (A2), we find:

$$\begin{aligned} u_{\mu\nu} \delta u^{\mu\nu} \sqrt{-g} &= u_{\mu\nu} \delta \left( g^{\mu\alpha} g^{\beta\nu} u_{\alpha\beta} \right) \sqrt{-g} \\ &= u_{\mu\nu} \left[ g^{\mu\alpha} g^{\beta\nu} \delta u_{\alpha\beta} + g^{\mu\alpha} u_{\alpha\beta} \delta g^{\beta\nu} + \right. \\ &\quad \left. + g^{\beta\nu} u_{\alpha\beta} \delta g^{\mu\alpha} \right] \sqrt{-g} \\ &= u^{\alpha\beta} \delta u_{\alpha\beta} \sqrt{-g} + 2u_{\mu\nu} g^{\mu\alpha} u_{\alpha\beta} \sqrt{-g} \delta g^{\beta\nu} \\ &= u^{\alpha\beta} \delta u_{\alpha\beta} \sqrt{-g} - 2g^{\nu\beta} u_{\kappa\nu} u^{\kappa\alpha} \sqrt{-g} \delta g_{\alpha\beta}. \end{aligned}$$

Substituting this expression in (B1) gives the following:

$$\left. \begin{aligned} & \delta \left( u_{\mu\nu} u^{\mu\nu} \sqrt{-g} \right) = 2u^{\alpha\beta} \delta u_{\alpha\beta} \sqrt{-g} \\ & - 2g^{\nu\beta} u_{\kappa\nu} u^{\kappa\alpha} \sqrt{-g} \delta g_{\alpha\beta} \\ & + \frac{1}{2} u_{\mu\nu} u^{\mu\nu} g^{\alpha\beta} \sqrt{-g} \delta g_{\alpha\beta}. \end{aligned} \right\} \quad (B2)$$

We will denote by  $B^{\alpha\beta}$  the stress-energy tensor of the acceleration field:

$$B^{\alpha\beta} = \frac{c^2}{4\pi\eta} \left( -g^{\alpha\nu} u_{\kappa\nu} u^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} u_{\mu\nu} u^{\mu\nu} \right). \quad (B3)$$

Given that  $u_{\mu\nu} = \nabla_\mu u_\nu - \nabla_\nu u_\mu = \partial_\mu u_\nu - \partial_\nu u_\mu$ , using differentiation by parts, as well as the equation which is valid for the antisymmetric tensor:  $\partial_\alpha (u^{\alpha\beta} \sqrt{-g}) = \sqrt{-g} \nabla_\alpha u^{\alpha\beta}$ , for the term  $2u^{\alpha\beta} \delta u_{\alpha\beta} \sqrt{-g}$  in (B2), we obtain:

$$\begin{aligned} & 2u^{\alpha\beta} \delta u_{\alpha\beta} \sqrt{-g} \\ &= 2u^{\alpha\beta} \delta (\partial_\alpha u_\beta - \partial_\beta u_\alpha) \sqrt{-g} \\ &= 2u^{\alpha\beta} (\partial_\alpha \delta u_\beta - \partial_\beta \delta u_\alpha) \sqrt{-g} \\ &= 4u^{\alpha\beta} \sqrt{-g} \partial_\alpha \delta u_\beta \\ &= 4\partial_\alpha (u^{\alpha\beta} \sqrt{-g} \delta u_\beta) - 4\partial_\alpha (u^{\alpha\beta} \sqrt{-g}) \delta u_\beta \\ &= 4\partial_\alpha (u^{\alpha\beta} \sqrt{-g} \delta u_\beta) - 4\nabla_\alpha u^{\alpha\beta} \sqrt{-g} \delta u_\beta. \end{aligned}$$

The term  $4\partial_\alpha (u^{\alpha\beta} \sqrt{-g} \delta u_\beta)$  in the last expression is the divergence, and it can be neglected in the variation of the action function.

Substituting the remaining term in (B2) and then in (B1) and using (B3), we find:

$$\delta S_7 = \int \left( \frac{c}{4\pi\eta} \nabla_\alpha u^{\alpha\beta} \delta u_\beta - \frac{1}{2c} B^{\alpha\beta} \delta g_{\alpha\beta} \right) \sqrt{-g} d\Sigma.$$

### Appendix C. Variation of the eighth term in the action function

Action variation for the eighth term in (1) has the form:

$$\delta S_8 = -\frac{1}{c} \int \delta (\pi_\mu J^\mu \sqrt{-g}) d\Sigma.$$

Acting like in Appendix A, taking (A2) and (A3) into account, we find:

$$\begin{aligned} & \delta (\pi_\mu J^\mu \sqrt{-g}) \\ &= \pi_\mu \sqrt{-g} \delta J^\mu + \pi_\mu J^\mu \delta \sqrt{-g} + J^\mu \sqrt{-g} \delta \pi_\mu \\ &= \pi_\mu \partial_\sigma \left[ \sqrt{-g} (J^\sigma \xi^\mu - J^\mu \xi^\sigma) \right] \\ &+ \frac{1}{2} \pi_\mu J^\mu g^{\alpha\beta} \sqrt{-g} \delta g_{\alpha\beta} + J^\mu \sqrt{-g} \delta \pi_\mu. \end{aligned}$$

$$\begin{aligned}
& \pi_\mu \partial_\sigma \left[ \sqrt{-g} (J^\sigma \xi^\mu - J^\mu \xi^\sigma) \right] \\
&= \partial_\sigma \left[ \pi_\mu \sqrt{-g} (J^\sigma \xi^\mu - J^\mu \xi^\sigma) \right] \\
&\quad - \sqrt{-g} (J^\sigma \xi^\mu - J^\mu \xi^\sigma) \partial_\sigma \pi_\mu.
\end{aligned}$$

In action variation, the term with the divergence is insignificant; the second term is transformed further:

$$\begin{aligned}
& -\sqrt{-g} (J^\sigma \xi^\mu - J^\mu \xi^\sigma) \partial_\sigma \pi_\mu \\
&= -(\partial_\sigma \pi_\mu - \partial_\mu \pi_\sigma) J^\sigma \xi^\mu \sqrt{-g} \\
&= f_{\mu\sigma} J^\sigma \xi^\mu \sqrt{-g}.
\end{aligned}$$

As a result, the variation of the eighth term equals:

$$\delta S_8 = \int \left( -\frac{1}{c} f_{\beta\sigma} J^\sigma \xi^\beta - \frac{1}{c} J^\beta \delta \pi_\beta - \frac{1}{2c} \pi_\mu J^\mu g^{\alpha\beta} \delta g_{\alpha\beta} \right) \sqrt{-g} d\Sigma.$$

#### Appendix D. Variation of the ninth term in the action function

In the special case when  $\sigma$  is a constant, for the variation of the ninth term in (1), with regard to (A2), we have:

$$\begin{aligned}
\delta S_9 &= -\frac{c}{16\pi\sigma} \int \delta (f_{\mu\nu} f^{\mu\nu} \sqrt{-g}) d\Sigma, \\
&\left. \begin{aligned}
& \delta (f_{\mu\nu} f^{\mu\nu} \sqrt{-g}) \\
&= \delta (f_{\mu\nu} f^{\mu\nu}) \sqrt{-g} + f_{\mu\nu} f^{\mu\nu} \delta \sqrt{-g} \\
&= f_{\mu\nu} \delta f^{\mu\nu} \sqrt{-g} + f^{\mu\nu} \delta f_{\mu\nu} \sqrt{-g} \\
&\quad + \frac{1}{2} f_{\mu\nu} f^{\mu\nu} g^{\alpha\beta} \sqrt{-g} \delta g_{\alpha\beta}.
\end{aligned} \right\} \quad (D1)
\end{aligned}$$

Replacing  $f^{\mu\nu} = g^{\mu\alpha} g^{\beta\nu} f_{\alpha\beta}$  and using  $\delta g^{\beta\nu}$  in (A2), we transform the first term of the equation and then substitute it in (D1):

$$\begin{aligned}
& f_{\mu\nu} \delta f^{\mu\nu} \sqrt{-g} \\
&= f_{\mu\nu} \delta (g^{\mu\alpha} g^{\beta\nu} f_{\alpha\beta}) \sqrt{-g} \\
&= f_{\mu\nu} \left[ g^{\mu\alpha} g^{\beta\nu} \delta f_{\alpha\beta} + g^{\mu\alpha} f_{\alpha\beta} \delta g^{\beta\nu} \right] \sqrt{-g} \\
&= f^{\alpha\beta} \delta f_{\alpha\beta} \sqrt{-g} + 2 f_{\mu\nu} g^{\mu\alpha} f_{\alpha\beta} \sqrt{-g} \delta g^{\beta\nu} \\
&= f^{\alpha\beta} \delta f_{\alpha\beta} \sqrt{-g} - 2 g^{\nu\beta} f_{\kappa\nu} f^{\kappa\alpha} \sqrt{-g} \delta g_{\alpha\beta}. \\
&\delta (f_{\mu\nu} f^{\mu\nu} \sqrt{-g}) \\
&= 2 f^{\alpha\beta} \delta f_{\alpha\beta} \sqrt{-g} \\
&\quad - 2 g^{\nu\beta} f_{\kappa\nu} f^{\kappa\alpha} \sqrt{-g} \delta g_{\alpha\beta} \\
&\quad + \frac{1}{2} f_{\mu\nu} f^{\mu\nu} g^{\alpha\beta} \sqrt{-g} \delta g_{\alpha\beta}.
\end{aligned}$$

We will denote by  $P^{\alpha\beta}$  the stress-energy tensor of the pressure field:

$$P^{\alpha\beta} = \frac{c^2}{4\pi\sigma} \left( -g^{\alpha\nu} f_{\kappa\nu} f^{\kappa\beta} + \frac{1}{4} g^{\alpha\beta} f_{\mu\nu} f^{\mu\nu} \right). \quad (D2)$$

We will transform the term  $2 f^{\alpha\beta} \delta f_{\alpha\beta} \sqrt{-g}$ , given that  $f_{\mu\nu} = \partial_\mu \pi_\nu - \partial_\nu \pi_\mu$ , using differentiation by parts as well as equation  $\partial_\alpha (f^{\alpha\beta} \sqrt{-g}) = \sqrt{-g} \nabla_\alpha f^{\alpha\beta}$  which is valid for the antisymmetric tensor:

$$\begin{aligned}
& 2 f^{\alpha\beta} \delta f_{\alpha\beta} \sqrt{-g} \\
&= 2 f^{\alpha\beta} \delta (\partial_\alpha \pi_\beta - \partial_\beta \pi_\alpha) \sqrt{-g} \\
&= 2 f^{\alpha\beta} (\partial_\alpha \delta \pi_\beta - \partial_\beta \delta \pi_\alpha) \sqrt{-g} \\
&= 4 f^{\alpha\beta} \sqrt{-g} \partial_\alpha \delta \pi_\beta \\
&= 4 \partial_\alpha (f^{\alpha\beta} \sqrt{-g} \delta \pi_\beta) \\
&\quad - 4 \partial_\alpha (f^{\alpha\beta} \sqrt{-g}) \delta \pi_\beta \\
&= 4 \partial_\alpha (f^{\alpha\beta} \sqrt{-g} \delta \pi_\beta) \\
&\quad - 4 \nabla_\alpha f^{\alpha\beta} \sqrt{-g} \delta \pi_\beta.
\end{aligned}$$

In the latter equation, the term with the divergence can be neglected, since it does not contribute to the variation of the action function. Substituting the results in (D1), we find the required variation:

$$\delta S_9 = \int \left( \begin{array}{c} \frac{c}{4\pi\sigma} \nabla_\alpha f^{\alpha\beta} \delta\pi_\beta \\ -\frac{1}{2c} P^{\alpha\beta} \delta g_{\alpha\beta} \end{array} \right) \sqrt{-g} d\Sigma.$$

### Appendix E. Acceleration tensor and equations for the acceleration field

By definition, the acceleration field appears as a result of applying the four-rotor to the four-potential:

$$u_{\mu\nu} = \nabla_\mu u_\nu - \nabla_\nu u_\mu = \partial_\mu u_\nu - \partial_\nu u_\mu.$$

The tensor  $u_{\mu\nu}$  is antisymmetric and includes various components of accelerations. By its structure, this tensor is similar to the gravitational tensor  $\Phi_{\mu\nu}$  and the electromagnetic tensor  $F_{\mu\nu}$ , each of which consists of two vector components depending on the field potentials and the velocities of the field sources.

In order to better understand the physical meaning of the acceleration field, we will introduce the following notations:

$$\begin{aligned} u_{0i} &= \partial_0 u_i - \partial_i u_0 = \frac{1}{c} S_i, \\ u_{ij} &= \partial_i u_j - \partial_j u_i = -N_k, \end{aligned} \quad (E1)$$

where the indices  $i, j, k$  form triples of non-recurring numbers of the form 1,2,3 or 3,1,2 or 2,3,1; three-vectors  $\mathbf{S}$  and  $\mathbf{N}$  can be written by components:

$$\begin{aligned} \mathbf{S} &= S_i = (S_1, S_2, S_3) = (S_x, S_y, S_z); \\ \mathbf{N} &= N_i = (N_1, N_2, N_3) = (N_x, N_y, N_z). \end{aligned}$$

Then, the tensor  $u_{\mu\nu}$  can be represented as follows:

$$u_{\mu\nu} = \begin{pmatrix} 0 & \frac{S_x}{c} & \frac{S_y}{c} & \frac{S_z}{c} \\ -\frac{S_x}{c} & 0 & -N_z & N_y \\ -\frac{S_y}{c} & N_z & 0 & -N_x \\ -\frac{S_z}{c} & -N_y & N_x & 0 \end{pmatrix}. \quad (E2)$$

In order to simplify our further arguments, we will consider the case of the flat spacetime; i.e., Minkowski space or the spacetime of the special theory of relativity. The role of the metric tensor in this case is played by the tensor  $\eta^{\alpha\beta}$ , the non-zero components of which are  $\eta^{00}=1$ ,  $\eta^{11}=\eta^{22}=\eta^{33}=-1$ . With its help, we will raise the indices of the acceleration tensor:

$$u^{\alpha\beta} = \eta^{\alpha\mu} \eta^{\nu\beta} u_{\mu\nu} = \begin{pmatrix} 0 & \frac{S_x}{c} & \frac{S_y}{c} & \frac{S_z}{c} \\ \frac{S_x}{c} & 0 & -N_z & N_y \\ \frac{S_y}{c} & N_z & 0 & -N_x \\ \frac{S_z}{c} & -N_y & N_x & 0 \end{pmatrix}. \quad (E3)$$

We will expand the four-vector of the mass current:  $J^\mu = \rho_0 u^\mu = \rho_0 (\gamma c, \gamma \mathbf{v})$ , where

$$\gamma = \frac{1}{\sqrt{1-v^2/c^2}}. \text{ In equations (22) and (23), we}$$

can replace the covariant derivatives  $\nabla_\beta$  with the partial derivatives  $\partial_\beta$ . Now, with the help of the vectors  $\mathbf{S}$  and  $\mathbf{N}$ , these equations can be presented as follows:

$$\begin{aligned} \nabla \cdot \mathbf{S} &= 4\pi\eta\gamma\rho_0, \quad \nabla \times \mathbf{N} = \frac{1}{c^2} \frac{\partial \mathbf{S}}{\partial t} + \frac{4\pi\eta\gamma\rho_0 \mathbf{v}}{c^2}, \\ \nabla \cdot \mathbf{N} &= 0, \quad \nabla \times \mathbf{S} = -\frac{\partial \mathbf{N}}{\partial t}. \end{aligned} \quad (E4)$$

The equations (E4), obtained in the framework of the special theory of relativity for the case  $\eta = \text{const}$ , are similar in their form to Maxwell equations in electrodynamics.

If we multiply scalarly the second equation in (E4) by  $\mathbf{S}$ , multiply scalarly the fourth equation

by  $-\mathbf{N}$  and sum up the results, we will obtain the following:

$$-\nabla \cdot [\mathbf{S} \times \mathbf{N}] = \left. \begin{aligned} & \frac{1}{2c^2} \frac{\partial (S^2 + c^2 N^2)}{\partial t} \\ & + \frac{4\pi\eta\gamma\rho_0 \mathbf{v} \cdot \mathbf{S}}{c^2} \end{aligned} \right\} \quad (\text{E5})$$

Equation (E5) contains the Poynting's theorem applied to the acceleration field. The meaning of this differential equation is that if in a system the work is done to accelerate the particles, then the power of this work is associated with the divergence of the acceleration field's flux and the change in time of the energy associated with the acceleration field. The relation (E5), in a generally covariant form according to (33), can be written as follows:

$$\nabla_\beta B^{0\beta} = -u^{0\beta} J_\beta.$$

We will now substitute (E2) in (30) and write the scalar and vector relations for the components of the four-acceleration  $a_\beta = (a_0, a_i)$ :

$$\left. \begin{aligned} \rho_0 \frac{du_0}{d\tau} &= \rho_0 a_0 = -u_{0\sigma} J^\sigma = -\frac{\gamma\rho_0}{c} \mathbf{S} \cdot \mathbf{v}, \\ \rho_0 \frac{du_i}{d\tau} &= \rho_0 a_i = -u_{i\sigma} J^\sigma \\ &= \gamma\rho_0 (\mathbf{S} + [\mathbf{v} \times \mathbf{N}]). \end{aligned} \right\} \quad (\text{E6})$$

The components of the four-acceleration are obtained from these relations after canceling  $\rho_0$ . As we can see, both vectors  $\mathbf{S}$  and  $\mathbf{N}$  make contribution to the space component  $a_i$  of the four-acceleration, the vector  $\mathbf{S}$  has the dimension of an ordinary three-acceleration and the dimension of the vector  $\mathbf{N}$  is the same as that of the frequency.

If we take into account that the four-potential of the acceleration field  $u_\mu = \left( \frac{g}{c}, -\mathbf{U} \right)$  in the case of one particle can be regarded as the covariant four-velocity, then from (E1) in Minkowski space, it follows:

$$\mathbf{S} = -\nabla g - \frac{\partial \mathbf{U}}{\partial t} = -c^2 \nabla \gamma - \frac{\partial(\gamma \mathbf{v})}{\partial t},$$

$$\mathbf{N} = \nabla \times \mathbf{U} = \nabla \times (\gamma \mathbf{v}). \quad (\text{E7})$$

The vector  $\mathbf{S}$  is the acceleration field strength and the vector  $\mathbf{N}$  is a quantity similar in its meaning to the magnetic field induction in electrodynamics or to the torsion field in the covariant theory of gravitation (the gravitomagnetic field in the general theory of relativity). At the constant velocity  $\mathbf{v} = \text{const}$ , the vectors  $\mathbf{S}$  and  $\mathbf{N}$  vanish. If there are non-zero time derivatives or spatial gradients of the velocity, then the acceleration field with the components  $\mathbf{S}$  and  $\mathbf{N}$  and the acceleration tensor  $u_{\mu\nu}$  appear. In this case, it is possible to state that the non-zero tensor  $u_{\mu\nu}$  in the inertial reference frame leads to the corresponding inertia forces as the consequence of any acceleration of bodies relative to the chosen reference frame.

If we substitute the tensors from (E2) and (E3) into (B3), then the stress-energy tensor of the acceleration field  $B^{\alpha\beta}$  will be expressed through the vectors  $\mathbf{S}$  and  $\mathbf{N}$ . In particular, for the tensor invariant  $u_{\mu\nu} u^{\mu\nu}$  and the time components of the tensor  $B^{\alpha\beta}$ , we have:

$$u_{\mu\nu} u^{\mu\nu} = -\frac{2}{c^2} (S^2 - c^2 N^2),$$

$$B^{00} = \frac{1}{8\pi\eta} (S^2 + c^2 N^2), \quad B^{0i} = \frac{c}{4\pi\eta} [\mathbf{S} \times \mathbf{N}]. \quad (\text{E8})$$

The component  $B^{00}$ , after its integration over the volume in the Lorentz reference frame, determines the energy of the acceleration field in the given volume, and the vector

$$\mathbf{K} = c B^{0i} = \frac{c^2}{4\pi\eta} [\mathbf{S} \times \mathbf{N}]$$

is the density of the energy flux of the acceleration field. Therefore, to calculate the energy flux of the acceleration field, the vector  $\mathbf{K}$  should also be integrated over the volume.

## Appendix F. The pressure tensor and equations for the pressure field

The pressure tensor is built by antisymmetric differentiation of the four-potential  $\pi_\nu$ :

$$f_{\mu\nu} = \nabla_\mu \pi_\nu - \nabla_\nu \pi_\mu = \partial_\mu \pi_\nu - \partial_\nu \pi_\mu.$$

We will introduce the following notations:

$$\begin{aligned} f_{0i} &= \partial_0 \pi_i - \partial_i \pi_0 = \frac{1}{c} C_i, \\ f_{ij} &= \partial_i \pi_j - \partial_j \pi_i = -I_k, \end{aligned} \quad (\text{F1})$$

where the indices  $i, j, k$  form triples of non-recurring numbers of the form 1,2,3 or 3,1,2 or 2,3,1; the three-vectors  $\mathbf{C}$  and  $\mathbf{I}$  in the Cartesian coordinates have the components:

$$\mathbf{C} = C_i = (C_1, C_2, C_3) = (C_x, C_y, C_z);$$

$$\mathbf{I} = I_i = (I_1, I_2, I_3) = (I_x, I_y, I_z).$$

In the specified notations, the tensor  $f_{\mu\nu}$  can be represented by the components:

$$f_{\mu\nu} = \begin{pmatrix} 0 & \frac{C_x}{c} & \frac{C_y}{c} & \frac{C_z}{c} \\ -\frac{C_x}{c} & 0 & -I_z & I_y \\ -\frac{C_y}{c} & I_z & 0 & -I_x \\ -\frac{C_z}{c} & -I_y & I_x & 0 \end{pmatrix}. \quad (\text{F2})$$

In Minkowski space, the metric tensor does not depend on the coordinates and time and consists of zeros and ones. In such space, the components of the tensor  $f^{\mu\nu}$  repeat the components of the tensor  $f_{\mu\nu}$  and differ only in the signs of the time components:

$$f^{\mu\nu} = \begin{pmatrix} 0 & -\frac{C_x}{c} & -\frac{C_y}{c} & -\frac{C_z}{c} \\ \frac{C_x}{c} & 0 & -I_z & I_y \\ \frac{C_y}{c} & I_z & 0 & -I_x \\ \frac{C_z}{c} & -I_y & I_x & 0 \end{pmatrix}. \quad (\text{F3})$$

Substituting in equations (26) and (27) the covariant derivatives  $\nabla_\beta$  with the partial derivatives  $\partial_\beta$ , we can represent these equations in the form of four equations for the vectors  $\mathbf{C}$  and  $\mathbf{I}$ :

$$\begin{aligned} \nabla \cdot \mathbf{C} &= 4\pi\sigma\gamma\rho_0, \\ \nabla \times \mathbf{I} &= \frac{1}{c^2} \frac{\partial \mathbf{C}}{\partial t} + \frac{4\pi\sigma\gamma\rho_0 \mathbf{v}}{c^2}, \quad \nabla \cdot \mathbf{I} = 0, \\ \nabla \times \mathbf{C} &= -\frac{\partial \mathbf{I}}{\partial t}. \end{aligned} \quad (\text{F4})$$

We will remind that the equations (F4), obtained in the framework of the special theory of relativity, are valid for the case of  $\sigma = \text{const}$ . Similarly to (E5), we obtain the equation of local pressure energy conservation:

$$-\nabla \cdot [\mathbf{C} \times \mathbf{I}] = \frac{1}{2c^2} \frac{\partial (C^2 + c^2 I^2)}{\partial t} + \frac{4\pi\sigma\gamma\rho_0 \mathbf{v} \cdot \mathbf{C}}{c^2}.$$

This equation also follows from equation (34) and can be written with the help of the tensor  $P^{\alpha\beta}$  according to (D2) as follows:

$$\nabla_\beta P^{0\beta} = -f^{0\beta} J_\beta.$$

Tensor invariant  $f_{\mu\nu} f^{\mu\nu}$  and the time components of the tensor  $P^{\alpha\beta}$  are expressed with the help of (F2) and (F3) through the vectors  $\mathbf{C}$  and  $\mathbf{I}$ :

$$\begin{aligned} f_{\mu\nu} f^{\mu\nu} &= -\frac{2}{c^2} (C^2 - c^2 I^2), \\ P^{00} &= \frac{1}{8\pi\sigma} (C^2 + c^2 I^2), \quad P^{0i} = \frac{c}{4\pi\sigma} [\mathbf{C} \times \mathbf{I}]. \end{aligned} \quad (\text{F5})$$

The component  $P^{00}$  of the stress-energy tensor of pressure determines the pressure energy density inside the bodies, and the vector

$\mathbf{F} = c P^{0i} = \frac{c^2}{4\pi\sigma} [\mathbf{C} \times \mathbf{I}]$  defines the density of the pressure energy flux.

We will now estimate the quantity  $f_{i\sigma} J^\sigma$  with the index  $i = 1, 2, 3$ . According to (31), this quantity determines the contribution of the pressure field into the total density of the force acting on the particle. In view of (F2), it turns out that the density of the pressure force has two components:

$$f_{i\sigma} J^\sigma = -\gamma \rho_0 (\mathbf{C} + [\mathbf{v} \times \mathbf{I}]).$$

For comparison, the time component is the density of the pressure force capacity divided by the speed of light:

$$f_{0\sigma} J^\sigma = \frac{\gamma \rho_0 \mathbf{v} \cdot \mathbf{C}}{c}.$$

The vector  $\mathbf{C}$  has the dimension of acceleration and the vector  $\mathbf{I}$  has the dimension of frequency. These vectors, with the help of (F1) and the definition of the four-potential of the pressure field  $\pi_\mu = \frac{p_0}{\rho_0 c^2} u_\mu = \left( \frac{\wp}{c}, -\mathbf{\Pi} \right)$  in Minkowski space, can be written as follows:

$$\begin{aligned} \mathbf{C} &= -\nabla \wp - \frac{\partial \mathbf{\Pi}}{\partial t} = -\nabla \left( \frac{\gamma p_0}{\rho_0} \right) - \frac{\partial}{\partial t} \left( \frac{\gamma p_0 \mathbf{v}}{\rho_0 c^2} \right), \\ \mathbf{I} &= \nabla \times \mathbf{\Pi} = \nabla \times \left( \frac{\gamma p_0 \mathbf{v}}{\rho_0 c^2} \right), \end{aligned} \quad (\text{F6})$$

where  $u_\mu$  denotes the four-velocity,  $p_0$  is the pressure in the frame of reference associated with the particle,  $\gamma = \frac{1}{\sqrt{1-v^2/c^2}}$  is the Lorentz factor and  $\mathbf{v}$  is the particle velocity.

The vector  $\mathbf{I}$ , according to its properties, is similar to the magnetic induction vector, and the vector  $\mathbf{C}$  is similar to the electric field strength. Motionless particles do not create the vector  $\mathbf{I}$ , and for the vanishing of the vector  $\mathbf{C}$  it is also necessary that the relation  $\gamma p_0 / \rho_0$  would not depend on the coordinates. In this case, the contribution of the pressure field into the acceleration of the particles will be zero.

## References:

- [1] Misner, C.W., Thorne, K.S. and Wheeler, J.A., “Gravitation”, (W.H. Freeman, San Francisco, CA, 1973).
- [2] Poplawski, N.J., A Lagrangian description of interacting energy, arXiv:gr-qc/0608031v2.
- [3] Fedosin, S.G., Hadronic Journal, 35(1) (2012) 35.
- [4] Fedosin, S.G., Advances in Natural Science, 5(4) (2012) 55.  
doi: 10.3968/2Fj.ans.1715787020120504.2023.
- [5] Hilbert, D., *Göttinger Nachrichten*, Math.-Phys., K1 (1915) 395.
- [6] Weyl, H., “Raum, Zeit, Materie. Vorlesungen über allgemeine Relativitätstheorie”, (J. Springer, Berlin, 1919).
- [7] Fock, V.A., “The Theory of Space, Time and Gravitation”, (Pergamon Press, London, 1959).
- [8] Dirac, P.A.M., “General Theory of Relativity”, (John Wiley & Sons, New-York, 1975).
- [9] Pauli, W., “Theory of Relativity”, (Dover Publications, New York, 1981).
- [10] Born, M., *Annalen der Physik*, 333(3) (1909) 571.
- [11] Fedosin, S.G., “Fizicheskie teorii i beskonechnaia vlozhennost' materii”, (Perm, 2009).
- [12] Утияма, Р., “Теория относительности”, (Атомиздат, Москва, 1979).
- [13] Landau, L.D. and Lifshitz, E.M., “The Classical Theory of Fields”, Vol. 2, 4<sup>th</sup> ed. (Butterworth-Heinemann, 1975).
- [14] Einstein, A., *Annalen der Physik*, 354(7) (1916) 769. doi: 10.1002/andp.19163540702.